

1617 01-23-02
360 #1/2

PATENT
674523-2012

04400

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant(s) : Kingsman et al.
U.S. Appln. No. : 10/016,686
U.S. Filing Date : November 2, 2002
Title of Invention : ANTIBODIES

RECEIVED
NOV 06 2002
TECH CENTER 1600/2900

APR 01 2002

RECEIVED

745 Fifth Avenue
New York, NY 10151

EXPRESS MAIL

Mailing Label Number: EV00141811623
Date of Deposit: January 17, 2002

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" Service under 37 CFR 1.10 on the date indicated above and is addressed to the Assistant Commissioner. For Patents, Washington, DC 20231.

Charles Jackson
(Typed or printed name of person mailing paper or fee)

Charles Jackson
(Signature of person mailing paper or fee)

RECEIVED
MAR 19 2002
Technology Center 2600

CLAIM OF PRIORITY

Assistant Commissioner for Patents
Washington, D.C. 20231

Sir:

Applicants hereby claim priority under 35 U.S.C. §119 and for 120, from British application numbers GB003527.9, GB0005071.6 and International patent application numbers PCT/GB00/04317 and PCT/GB99/03859, a certified copy of each is enclosed.

Acknowledgment of the claim of priority and of the receipt of said certified copy are respectfully requested.

Respectfully submitted,

FROMMER LAWRENCE & HAUG LLP

By: Thomas J. Kowalski
THOMAS J. KOWALSKI, Reg. No. 32,147
(212) 588-0800

THIS PAGE BLANK (USPTO)



TECH CENTER 1600/2900

RECEIVED
NOV 06 2002

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

INVESTOR IN PEOPLE
APR 01 2002
TECH CENTER 1600/2900
RECEIVED

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.


Signed

Stephen Horvath

Dated 6 December 2001

THIS PAGE BLANK (USPTO)





16FEB00 E513691-8 D02246
P01/7700 0.00-0003527.9

Request for a grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

The Patent Office

Cardiff Road
Newport
Gwent NP9 1RH

1.	Your reference	P008161GB CTH		
2.	Patent application number (The Patent Office will fill in this part)	0003527.9	15 FEB 2000	
3.	Full name, address and postcode of the or of each applicant (underline all surnames)	Oxford BioMedica (UK) Limited Medawar Centre Robert Robinson Avenue The Oxford Science Park Oxford OX4 4GA Patents ADP number (if you know it) 7667157001 If the applicant is a corporate body, give the country/state of its incorporation United Kingdom		
4.	Title of the invention	Antibodies		
5.	Name of your agent (if you have one)	D YOUNG & CO		
	"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)	21 NEW FETTER LANE LONDON EC4A 1DA Patents ADP number (if you have one) 59006		
6.	If you are declaring priority from one or more earlier patent applications, give the country and date of filing of the or each of these earlier applications and (if you know it) the or each application number	Country	Priority application number (if you know it)	Date of filing (day/month/year)
		PCT	PCT/GB99/03859	18/11/99
7.	If this application is divided or otherwise derived from an earlier UK application, give the number and filing date of the earlier application	Number of earlier application	Date of filing (day/month/year)	

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:
a) any applicant named in part 3 is not an inventor, or
b) there is an inventor who is not named as an applicant, or
c) any named applicant is a corporate body.
See note (d))

Yes

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form 0

Description 78 + 1 (SEQUENCE LISTING)

Claims(s) 6

Abstract 1

Drawing(s) 20 M

10. If you are also filing any of the following, state how many against each item.

Priority documents 0

Translations of priority documents 0

Statement of inventorship and right to grant of a patent (Patents Form 7/77) 0

Request for preliminary examination and search (Patents Form 9/77) 0

Request for substantive examination (Patents Form 10/77) 0

Any other documents 0
(please specify)

11. I/We request the grant of a patent on the basis of this application.

Signature

Date


D. YOUNG & CO
Agents for the Applicants

15 02 00

12. Name and daytime telephone number of the person to contact in the United Kingdom

~~Catherine Maffrand~~

023 8063 4816

C. T. HARDING

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

a) If you need help to fill in this form or you have any questions, please contact the Patent Office on 01645 500505

b) Write your answers in capital letters using black ink or you may type them

c) If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.

d) If you answered 'Yes' Patents Form 7/77 will need to be filed.

e) Once you have filled in the form you must remember to sign and date it.

f) For details of the fee and ways to pay please contact the Patent Office.

ANTIBODIES

FIELD OF THE INVENTION

5

The present invention relates to antibodies.

In particular, the present invention relates to antibodies that recognise a disease associated cell surface marker (DACSM).

10

More particularly, the present invention relates *in vitro* and *in vivo/ex vivo* applications of these antibodies in the diagnosis and treatment of diseases associated with a DACSM.

15 BACKGROUND TO THE INVENTION

In certain disease states, a derangement of cellular metabolism can affect the level of expression of one or more DACSMs. In some circumstances, this cellular derangement may lead to a change in the levels of expression of the DACSM. Thus, each disease
20 causing agent or disease state may have associated with it a DACSM which may be crucial in the immune recognition and/or the elimination and/or control of a disease causing agent or disease state in a host organism. In this way, the DACSM may be capable of acting as a marker not only for the diagnosis of disease states but also for the accurate staging of the disease profile so that the appropriate therapy may be designed.

25

A particular example of DACSMs which have been well characterised include the tumour-associated antigens (TAAs). A number of oncofoetal or tumour-associated antigens (TAAs) have been identified and characterised in human and animal tumours. These TAAs include carcinoembryonic antigen (CEA), TAG72, c-erbB2,
30 (underglycosylated) MUC-1 and p53, epithelial glycoprotein-2 antigen (EGP-2; also known as EGP40, Ep-CAM, KSA, CO17-1A or GA733-2) and the 5T4 antigen. In general, TAAs are antigens which are expressed during foetal development but which

are downregulated in adult cells, and are thus normally absent or present only at very low levels in adults. However, during tumourigenesis, tumour cells have been observed to resume expression of TAAs. Thus, it is thought that malignant cells may be distinguished from their non-malignant counterparts by resumption of expression of TAAs. Consequently, application of TAAs for (i) *in vitro* and/or *in vivo/ex vivo* diagnosis of tumour disorders; (ii) for imaging and/or immunotherapy of cancer has been suggested and (iii) as indicators of progression of tumour associated disease;

In order to mount a humoral and/or cellular immune response against a particular disease, the host immune system must come in contact with a DACSM. In addition to recognising foreign antigens, T cells often need additional stimulation to become fully activated. It is now becoming apparent that two signals are required for activation of naive T-cells by antigen bearing target cells. One signal is an antigen specific signal, delivered through the T-cell receptor and the second signal is an antigen independent or co-stimulatory signal leading to lymphokine products. These additional signals are delivered through other receptors (such as CD28 and CD40L) on the T cell that interact with ligands (such as B7 and CD40) which are present on professional antigen presenting cells (APCs), such as dendritic cells and macrophages, but which are absent from other cells. These co-stimulatory ligands are often referred to as co-stimulatory molecules.

By way of example, the B7 family (namely B7.1, B7.2, and possibly B7.3) represent a recently discovered, but important group of co-stimulatory molecules. B7.1 and B7.2 are both member of the Ig gene superfamily. If a T lymphocyte encounters an antigen alone, without co-stimulation by B7, it will respond with either anergy, or apoptosis (programmed cell death). If the co-stimulatory signal is provided it will respond with clonal expansion against the target antigen. No significant amplification of the immune response against a given antigen is thought to occur without co-stimulation (June *et al* (Immunology Today 15:321-331, 1994); Chen *et al* (Immunology Today 14:483-486); Townsend *et al* (Science 259:368-370)). Freeman *et al* (J. Immunol. 143:2714-2722, 1989). Azuma *et al* (Nature 366:76-79, 1993). Thus, it has been postulated that one method for stimulating immune recognition of diseased cells which are poorly

immunogenic would be to enhance antigen presentation and co-stimulation of lymphocytes in the presence of the DACSM.

By way of example, it has been shown that disease states such as cancer, established
5 tumours may be poorly immunogenic despite the fact that they commonly express
DACSMs. Transfection of the genes encoding B7-1 and B7-2, either alone or in
combination with cytokines, have been shown to enhance the development of immunity
to experimental tumours in animal models (e.g. Leong *et al.* 1997 *Int. J. Cancer* 71:
476-482; Zitvogel *et al.* 1996 *Eur. J. Immunol.* 26:1335-1341; Cayeux *et al.* 1997 *J.*
10 *Immunol* 158:2834-2841). However, in translating these results into a practical
treatment for human cancer, there are a number of significant problems to be overcome.
A major problem in such studies has been the need to deliver B7 genes *in vivo* to a large
number of cells of the tumour to achieve efficacy. A second problem has been the
selective target expression of B7 to the tumour cells to avoid inappropriate immune cell
15 activation directed against other cell types. Some solutions to these problems have been
addressed in WO 98/55607 where a tumour interacting protein (TIP) such as a tumour
binding protein (TBP) has been used to selectively target a co-stimulatory molecule to
tumour cells.

20 Recombinant DNA technologies have been applied to develop antibodies that recognise
DACSMs (Hoogenboom *et al* 1998 *Immunotechnology* 4: 1-20; and Winter 1998 *FEBS*
Lett 458: 92-94. Recently, there has been considerable interest in using antibody gene
libraries to generating antibodies, such as a single chain antibody (ScFv Abs). It is well
known that in certain circumstances, there are advantages of using ScFv Abs, rather
25 than whole antibodies. The smaller size of the fragments allows for rapid clearance,
and may lead to improved tumour to non-tumour ratios. However, many efforts have
failed to produce ScFv Abs of high specificity. Moreover, whole IgGs are regarded as a
better format for therapeutic Mabs than ScFc Abs as they are regarded as having an
extended serum half life (see Vaughan *et al* 1998, *Nature Biotech* 16: 535-539).

30

The present invention seeks to provide an ScFv Ab raised against a DACSM which is
useful in the treatment of disease conditions associated with a DACSM.

SUMMARY ASPECTS OF THE PRESENT INVENTION

The present invention provides an ScFv Ab (ScFv Ab), capable of recognising a DACSM and having a therapeutic effect in diseases associated with a DACSM. This ScFv Ab can be directly administered either as a peptide (synthetically or genetically expressed) or as "naked DNA" (for example, in a plasmid) or via a delivery vehicle such as a viral vector comprising the nucleotide sequence encoding the ScFv Ab. For some cases, this ScFv Ab may be more efficacious than a ScFv Ab fused to an secreted co-stimulatory molecule (SCM) such as B7 or IgG. Using an ScFv Ab was not an obvious choice as a therapeutic agent, for the treatment of diseases such as cancer, especially as one would expect that a fusion protein comprising a SCM fused to an ScFv would perform better than an ScFv alone.

The present invention is advantageous for the following reasons:

- (i) it provides an ScFv Ab capable of recognising a DACSM. For some cases, it has a greater therapeutic effect than an ScFv Ab which is fused to a SCM such as B7 or an immunoglobulin such as IgG;
- (ii) it provides a high affinity ScFv Ab which has applications in:
 - (a) *in vitro* and *in vivo/ex vivo* diagnosis and therapy;
 - (b) imaging and the treatment of cells expressing the DACSM antigen;
 - (c) prevention and/or treatment of different human diseases such as carcinomas when the ScFv Ab is used either alone or in combination with suitable diagnostic and/or therapeutically useful agents;
 - (d) studies relating to the isolation and/or purification of the DACSM antigen to which the ScFv Abs specifically binds; and

(e) providing building blocks for further rational therapeutic ScFv Ab design and screens for ScFv Abs capable of binding to target DACSMs and/or screens for DACSMs capable of binding to target ScFv Abs.

5 DETAILED ASPECTS OF THE INVENTION

Other aspects of the present invention are presented in the accompanying claims and in the following description and drawings. These aspects are presented under separate section headings. However, it is to be understood that the teachings under each section
10 are not necessarily limited to that particular section heading.

ScFv ANTIBODY

In one aspect, the present invention provides a recombinant ScFv Ab that recognises a
15 DACSM antigen.

As used herein, the term "ScFv Ab" means an antibody capable of recognising a DACSM antigen which has a light chain variable region (VL) and a heavy chain variable (VH) region. The VH and VL partner domains are typically linked/joined via a
20 flexible oligopeptide/peptide linker. The VH and VL partner domains may be connected in the order of VH followed by VL or VL followed by VH. Typically, the the sequences may be connected via a linker sequence in the order VH-linker-VL or VL-linker-VH. As used herein, the term includes fragments of proteolytically-cleaved or recombinantly-prepared portions of an ScFv Ab molecule that are capable of
25 selectively reacting with or recognising a DACSM. Non limiting examples of such proteolytic and/or recombinant fragments include chimeric ScFv antibodies which, for the purposes of this invention, may refer to an ScFv Ab having either a or both heavy and light chain variable regions (VH and VL) encoded by a nucleotide sequence derivable from a mammalian immunoglobulin gene other than a human
30 immunoglobulin gene and either a or both heavy and light chain encoded by a nucleotide sequence derivable from a human immunoglobulin gene. The ScFv Ab may be covalently or non-covalently linked to another entity (such as another ScFv Ab) to

form antibodies having two or more binding sites. For example, one ScFv Ab could bind to a DACSM, such as 5T4, and the second ScFv Ab could bind to an immune enhancer molecule.

5 In accordance with the present invention, reference to the term "ScFv Ab" includes but is not limited to reference to the peptide *per se* also as well the peptide as part of a fusion protein as well as the nucleotide sequence encoding the peptide and/or the nucleotide sequence encoding the fusion protein. The peptide *per se* and/or fusion protein may be a synthetic peptide. Alternatively, the peptide and/or fusion protein may
10 be a genetically expressed/recombinant peptide/fusion protein. For some applications, the term "ScFv Ab means peptide *per se*. The term "ScFv Ab" also includes an ScFv Ab with a secretion leader (L) sequence which is designated herein as LScFv.

As used herein, the term "variable region" refers to the variable region, or domain, of
15 the light chain (VL) and heavy chain (VH) which contain the determinants for binding recognition specificity and for the overall affinity of the ScFv Ab for a DACSM. The variable domains of each pair of light (VL) and heavy chains (VH) are involved in antigen recognition and form the antigen binding site. The domains of the light and heavy chains have the same general structure and each domain has four framework
20 (FR) regions, whose sequences are relatively conserved, connected by three complementarity determining regions (CDRs). The FR regions maintain the structural integrity of the variable domain. The CDRs are the polypeptide segments within the variable domain that mediate binding of an antigen such as a DACSM.

25 Preferably the affinity (K_D) of the ScFv Ab of the present invention for the 5T4 antigen is from about 5×10^{-10} to about 10×10^{-10} .

Preferably the affinity (K_D) of the ScFv Ab of the present invention for the 5T4 antigen is from about 6×10^{-10} to about 9×10^{-10} .

30

Preferably the affinity (K_D) of the ScFv Ab of the present invention for the 5T4 antigen is from about 7×10^{-10} to about 8×10^{-10} .

Preferably the affinity (K_D) of the ScFv Ab of the present invention for the 5T4 antigen is about 7.9×10^{-10} . The K_D of the ScFvAb is measured using BIAevaluation software (Pharmacia).

5 As used herein, the term "off-rate" means the dissociation rate (k_{off}) of a ScFv Ab from an antigen. In the context of the present invention, it is measured using BIAevaluation software (Pharmacia). A low off rate is desirable as it reflects the affinity of an Fab fragment for an antigen such as a DACSM.

10 As used herein, the term "affinity" is defined in terms of the dissociation rate or off-rate (k_{off}) of a ScFv Ab from a DACSM antigen. The lower the off-rate the higher the affinity that a ScFv Ab has for an antigen such as a DACSM.

DACSM

15

In accordance with the present invention the DACSM can include but is not limited to a receptor for adhesive proteins such as growth factor receptors. Examples of growth factor receptors include but are not limited to ApoE, Apo-SAA, BDNF, Cardiotrophin-1, EGF, ENA-78, Eotaxin, Eotaxin-2, Exodus-2, FGF-acidic, FGF-basic, fibroblast growth factor-10 (Marshall 1998 Nature Biotechnology 16: 129).FLT3 ligand (Kimura *et al* (1997), Fractalkine (CX3C), GDNF, G-CSF, GM-CSF, GF- β 1, insulin, IFN- γ , IGF-I, IGF-II, IL-1 α , IL-1 β , IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8 (72 a.a.), IL-8 (77 a.a.), IL-9, IL-10, IL-11, IL-12, IL-13, IL-15, IL-16, IL-17, IL-18 (IGIF), Inhibin α , Inhibin β , IP-10, keratinocyte growth factor-2 (KGF-2), KGF, Leptin, LIF, Lymphotoctin, Mullerian inhibitory substance, monocyte colony inhibitory factor, monocyte attractant protein (Marshall 1998 *ibid*), M-CSF, MDC (67 a.a.), MDC (69 a.a.), MCP-1 (MCAF), MCP-2, MCP-3, MCP-4, MDC (67 a.a.), MDC (69 a.a.), MIG, MIP-1 α , MIP-1 β , MIP-3 α , MIP-3 β , MIP-4, myeloid progenitor inhibitor factor-1 (MPIF-1), NAP-2, Neurturin, Nerve growth factor, β -NGF, NT-3, NT-4, Oncostatin M, PDGF-AA, PDGF-AB, PDGF-BB, PF-4, RANTES, SDF1 α , SDF1 β , SCF, SCGF, stem cell factor (SCF), TARC, TGF- α , TGF- β , TGF- β 2, TGF- β 3, tumour necrosis factor (TNF), TNF- α , TNF- β , TNIL-1, TPO, VEGF, GCP-2, GRO/MGSA, GRO- β , GRO- γ ,

20
25
30

HCC1, 1-309. A non-exhaustive list of growth factor receptors can be found on pages 392-297 Molecular Biology and Biotechnology (Ed RA Meyers 1995 VCH Publishers Inc.); a plasminogen activator; a metalloproteinase (such as collagenase), a mucin; a glycoprotein; an antigen restricted in its tissue distribution; and/or a cell surface molecule which plays a role in tumour cell growth, migration or metastasis, (such as a 5T4 antigen, a tumour specific carbohydrate moiety or an oncofetal antigen). The term DACSM may also includes antigenic determinants.

ANTIGENIC DETERMINANT

10

As used herein, the term "antigenic determinant" refers to any antigen which is associated with a disease or a disorder. By way of example, the antigenic determinant may also be derived from pathogenic agents associated with diseased cells, such as tumour cells, which multiply unrestrictedly in an organism and may thus lead to pathological growths. Examples of such pathogenic agents are described in Davis, B.D. *et al* (Microbiology, 3rd ed., Harper International Edition). The antigenic determinant may be an antigen and/or an immunodominant epitope on an antigen. By way of example, the antigenic determinant may include tumour associated antigens (TAA) which may serve as targets for the host immune system and elicit responses which result in tumour destruction.

20

TAA

The term "tumour associated antigen (TAA)" is used herein to refer to any TAA or antigenic peptide thereof. The antigen being one that is expressed by the tumour itself or cells associated with the tumour such as parenchymal cells or those of the associated vasculature. The term "tumour associated antigen (TAA)" includes antigens that distinguish the tumour cells from their normal cellular counterparts where they may be present in trace amounts.

30

Examples of TAAs include but are not limited to MART-1 (Melanoma Antigen Recognised by T cells-1) MAGE-1, MAGE-3, 5T4, gp100, Carcinoembryonic antigen

(CEA), prostate-specific antigen (PSA), MUCIN (MUC-1), tyrosinase. Particularly preferred TAAs are cell surface molecules as these are positioned for recognition by elements of the immune system and are excellent targets for therapy such as therapy and/or immunotherapy. The present invention is in no way limited to antigenic determinants encoding the above listed TAAs. Other TAAs may be identified, isolated and cloned by methods known in the art such as those disclosed in U.S. Patent No. 4,514,506.

5T4 TAA

10

The TAA 5T4 (see WO 89/07947) has been extensively characterised. It is a 72kDa glycoprotein expressed widely in carcinomas, but having a highly restricted expression pattern in normal adult tissues. It appears to be strongly correlated to metastasis in colorectal and gastric cancer. The full nucleic acid sequence of human 5T4 is known (Myers *et al.*, 1994 J Biol Chem 169: 9319-24).

15

CO-STIMULATORY MOLECULES

In order to respond to a DACSM, lymphocytes require at least two distinct signals to activate their effector functions (Bretscher and Cohn 1970 Science 169: 1042-1049; Crabtree 1989 Science 243: 355-361). The primary signal is specific for antigen. Stimulation of the primary signal in isolation normally leads to apoptosis (programmed cell death) of the lymphocyte or leads to the establishment of a state of sustained unresponsiveness or anergy (Weiss *et al. supra*). In order to achieve activation of the lymphocyte, accessory signals are required which may be delivered by cytokines or by cell-surface co-stimulatory ligands present on antigen-presenting cells (APC).

25

There are a number of such co-stimulatory molecules now identified including adhesion molecules, LFA-3, ICAM-1, ICAM-2. Major co-stimulatory molecules present on APC are the members of the B7 family including B7-1 (CD80), B7-2 (CD86) and B7-3. These molecules are ligands of co-stimulatory receptors on lymphocytes including CD28 (WO92/00092), probably the most significant co-stimulatory receptor for resting

30

T-cells. Different members of the B7 family of glycoproteins may deliver subtly different signals to T-cells (Nunes *et al.* 1996 J. Biol. Chem. 271: 1591-1598).

5 In one embodiment of the present invention, an ScFv Ab is used which comprises a secreted co-stimulatory molecule ("SCM") with binding affinity for a DACSM, such as a tumour antigen.

ScFv Ab SOURCE

10 The ScFv Ab of the present invention is obtainable from or produced by any suitable source, whether natural or not, or it may be a synthetic ScFv Ab, a semi-synthetic ScFv Ab, a mimetic, a derivatised ScFv Ab, a recombinant ScFv Ab, a fermentation optimised ScFv Ab, a fusion protein or equivalents, mutants and derivatives thereof as long as it retains the required DACSM binding specificity of the ScFv Ab of the present
15 invention. These include a ScFv Ab with DACSM binding specificity which may have amino acid substitutions or may have sugars or other molecules attached to amino acid functional groups.

The term "mimetic" relates to any chemical which may be a peptide, polypeptide,
20 antibody or other organic chemical which has the same binding specificity as the ScFv Ab of the present invention.

The term "derivative" or "derivatised" as used herein includes chemical modification of an ScFv Ab. Illustrative of such modifications would be replacement of hydrogen by
25 an alkyl, acyl, or amino group. Preferably, the ScFv Ab includes at least a portion of which has been prepared by recombinant DNA techniques or produced by chemical synthesis techniques or combinations thereof.

Preferably, the ScFv Ab is prepared by the use of chemical synthesis techniques.
30

CHEMICAL SYNTHESIS METHODS

The ScFv Ab of the present invention or variants, homologues, derivatives, fragments or mimetics thereof may be produced using chemical methods to synthesize the ScFv Ab amino acid sequence, in whole or in part. For example, peptides can be synthesized by solid phase techniques, cleaved from the resin, and purified by preparative high performance liquid chromatography (e.g., Creighton (1983) *Proteins Structures And Molecular Principles*, WH Freeman and Co, New York NY). The composition of the synthetic peptides may be confirmed by amino acid analysis or sequencing (e.g., the Edman degradation procedure; Creighton, *supra*).

Direct synthesis of the ScFv Ab or variants, homologues, derivatives, fragments or mimetics thereof can be performed using various solid-phase techniques (Roberge JY *et al* (1995) *Science* 269: 202-204) and automated synthesis may be achieved, for example, using the ABI 431A Peptide Synthesizer (Perkin Elmer) in accordance with the instructions provided by the manufacturer. Additionally, the amino acid sequences obtainable from the ScFv Ab, or any part thereof, may be altered during direct synthesis and/or combined using chemical methods with a sequence from other subunits, or any part thereof, to produce a variant ScFv Ab.

In an alternative embodiment of the invention, the coding sequence of the ScFv Ab or variants, homologues, derivatives, fragments or mimetics thereof may be synthesized, in whole or in part, using chemical methods well known in the art (see Caruthers MH *et al* (1980) *Nuc Acids Res Symp Ser* 215-23, Horn T *et al* (1980) *Nuc Acids Res Symp Ser* 225-232).

Preferably the ScFv Ab of the present invention comprises the amino acid sequence set out in SEQ ID No 1 (see Figure 1).

Preferably the ScFv Ab of the present invention comprises the amino acid sequence set out in SEQ ID No 3 (see Figure 2).

Preferably the ScFv Ab of the present invention comprises the amino acid sequence set out in SEQ ID No 4 (see Figure 6).

AMINO ACID SEQUENCES

5

As used herein, the term "amino acid sequence" refers to peptide, polypeptide sequences, protein sequences or portions thereof.

10

Preferably, the ScFv Ab is an isolated ScFv Ab and/or purified and/or non-native ScFv Ab.

15

The ScFv Ab of the present invention may be in a substantially isolated form. It will be understood that the protein may be mixed with carriers or diluents which will not interfere with the intended purpose of the ScFv Ab and still be regarded as substantially isolated. The ScFv Ab of the present invention may also be in a substantially purified form, in which case it will generally comprise the ScFv Ab in a preparation in which more than 90%, e.g. 95%, 98% or 99% of the ScFv Ab in the preparation is a peptide comprising SEQ ID No 1 or SEQ ID No 3 or SEQ ID No 4 or variants, homologues, derivatives or fragments thereof.

20

VARIANTS /HOMOLOGUES/DERIVATIVES OF AMINO ACID SEQUENCES

25

Preferred amino acid sequences of the present invention are set out in SEQ ID No 1 or SEQ ID No 3 or SEQ ID No 4 are sequences obtainable from the ScFv Ab of the present invention but also include homologous sequences obtained from any source, for example related viral/bacterial proteins, cellular homologues and synthetic peptides, as well as variants or derivatives thereof.

30

Thus, the present invention covers variants, homologues or derivatives of the amino acid sequences presented herein, as well as variants, homologues or derivatives of the nucleotide sequence coding for those amino acid sequences.

In the context of the present invention, a homologous sequence is taken to include an amino acid sequence which is at least 75, 85 or 90% identical, preferably at least 95 or 98% identical at the amino acid level over at least, for example, the amino acid sequence as set out in SEQ ID No 1 or SEQ ID No 3 or SEQ ID No 4 of the sequence listing herein. In particular, homology should typically be considered with respect to those regions of the sequence known to be essential for binding specificity (such as amino acids at positions) rather than non-essential neighbouring sequences. Although homology can also be considered in terms of similarity (i.e. amino acid residues having similar chemical properties/functions), in the context of the present invention it is preferred to express homology in terms of sequence identity.

Homology comparisons can be conducted by eye, or more usually, with the aid of readily available sequence comparison programs. These commercially available computer programs can calculate % homology between two or more sequences.

% homology may be calculated over contiguous sequences, i.e. one sequence is aligned with the other sequence and each amino acid in one sequence is directly compared with the corresponding amino acid in the other sequence, one residue at a time. This is called an "ungapped" alignment. Typically, such ungapped alignments are performed only over a relatively short number of residues.

Although this is a very simple and consistent method, it fails to take into consideration that, for example, in an otherwise identical pair of sequences, one insertion or deletion will cause the following amino acid residues to be put out of alignment, thus potentially resulting in a large reduction in % homology when a global alignment is performed. Consequently, most sequence comparison methods are designed to produce optimal alignments that take into consideration possible insertions and deletions without penalising unduly the overall homology score. This is achieved by inserting "gaps" in the sequence alignment to try to maximise local homology.

However, these more complex methods assign "gap penalties" to each gap that occurs in the alignment so that, for the same number of identical amino acids, a sequence alignment

with as few gaps as possible - reflecting higher relatedness between the two compared sequences - will achieve a higher score than one with many gaps. "Affine gap costs" are typically used that charge a relatively high cost for the existence of a gap and a smaller penalty for each subsequent residue in the gap. This is the most commonly used gap scoring system. High gap penalties will of course produce optimised alignments with fewer gaps. Most alignment programs allow the gap penalties to be modified. However, it is preferred to use the default values when using such software for sequence comparisons. For example when using the GCG Wisconsin Bestfit package (see below) the default gap penalty for amino acid sequences is -12 for a gap and -4 for each extension.

10

Calculation of maximum % homology therefore firstly requires the production of an optimal alignment, taking into consideration gap penalties. A suitable computer program for carrying out such an alignment is the GCG Wisconsin Bestfit package (University of Wisconsin, U.S.A.; Devereux *et al.*, 1984, *Nucleic Acids Research* 12:387). Examples of other software than can perform sequence comparisons include, but are not limited to, the BLAST package (see Ausubel *et al.*, 1999 *ibid* – Chapter 18), FASTA (Atschul *et al.*, 1990, *J. Mol. Biol.*, 403-410) and the GENWORKS suite of comparison tools. Both BLAST and FASTA are available for offline and online searching (see Ausubel *et al.*, 1999 *ibid*, pages 7-58 to 7-60). However it is preferred to use the GCG Bestfit program. A new tool, called BLAST 2 Sequences is also available for comparing protein and nucleotide sequence (see *FEMS Microbiol Lett* 1999 174(2): 247-50; *FEMS Microbiol Lett* 1999 177(1): 187-8 and tatiana@ncbi.nlm.nih.gov).

20

Although the final % homology can be measured in terms of identity, the alignment process itself is typically not based on an all-or-nothing pair comparison. Instead, a scaled similarity score matrix is generally used that assigns scores to each pairwise comparison based on chemical similarity or evolutionary distance. An example of such a matrix commonly used is the BLOSUM62 matrix - the default matrix for the BLAST suite of programs. GCG Wisconsin programs generally use either the public default values or a custom symbol comparison table if supplied (see user manual for further details). It is preferred to use the public default values for the GCG package, or in the case of other software, the default matrix, such as BLOSUM62.

30

Once the software has produced an optimal alignment, it is possible to calculate % homology, preferably % sequence identity. The software typically does this as part of the sequence comparison and generates a numerical result.

- 5 The terms "variant" or "derivative" in relation to the amino acid sequences of the present invention includes any substitution of, variation of, modification of, replacement of, deletion of or addition of one (or more) amino acids from or to the sequence providing the resultant amino acid sequence has a binding specificity, preferably having at least the same binding specificity as the amino acid sequence set out in SEQ ID No 1 or SEQ ID No 3 or
10 SEQ ID No 4 of the sequence listing herein.

SEQ ID No 1 or SEQ ID No 3 or SEQ ID No 4 of the sequence listing herein may be modified for use in the present invention. Typically, modifications are made that maintain the binding specificity of the sequence. Amino acid substitutions may be
15 made, for example from 1, 2 or 3 to 10 or 20 substitutions provided that the modified sequence retains the required binding specificity. Amino acid substitutions may include the use of non-naturally occurring analogues.

The ScFv Ab of the present invention may also have deletions, insertions or
20 substitutions of amino acid residues which produce a silent change and result in a functionally equivalent ScFv Ab. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues as long as the binding specificity of the ScFv Ab is retained. For example, negatively charged amino acids include aspartic acid
25 and glutamic acid; positively charged amino acids include lysine and arginine; and amino acids with uncharged polar head groups having similar hydrophilicity values include leucine, isoleucine, valine, glycine, alanine, asparagine, glutamine, serine, threonine, phenylalanine, and tyrosine.

30 Conservative substitutions may be made, for example according to the Table below. Amino acids in the same block in the second column and preferably in the same line in the third column may be substituted for each other:

ALIPHATIC	Non-polar	G A P
		I L V
	Polar - uncharged	C S T M
		N Q
	Polar - charged	D E
		K R
AROMATIC		H F W Y

Preferably, the isolated ScFv Ab and/or purified ScFv Ab and/or non-native ScFv Ab is prepared by use of recombinant techniques.

5

NUCLEOTIDE SEQUENCES

It will be understood by a skilled person that numerous different nucleotide sequences can encode the same ScFv Ab of the present invention as a result of the degeneracy of the genetic code. In addition, it is to be understood that skilled persons may, using routine techniques, make nucleotide substitutions that do not affect the ScFv Ab encoded by the nucleotide sequence of the present invention to reflect the codon usage of any particular host organism in which the ScFv Ab of the present invention is to be expressed.

The terms "variant", "homologue" or "derivative" in relation to the nucleotide sequence set out in SEQ ID No 5 (see Figure 1) or SEQ ID No 7 (see Figure 2) or SEQ ID No 8 (see Figure 6) of the present invention includes any substitution of, variation of, modification of, replacement of, deletion of or addition of one (or more) nucleic acid from or to the sequence providing the resultant nucleotide sequence codes for a ScFv Ab having a binding specificity, preferably having at least the same binding specificity as the nucleotide sequence set out in SEQ ID No 5 or SEQ ID No 7 or SEQ ID No 8 of the sequence listings of the present invention.

20

As indicated above, with respect to sequence homology, preferably there is at least 75%, more preferably at least 85%, more preferably at least 90% homology to the sequences shown in the sequence listing herein. More preferably there is at least 95%, more preferably at least 98%, homology. Nucleotide homology comparisons may be conducted
5 as described above. A preferred sequence comparison program is the GCG Wisconsin Bestfit program described above. The default scoring matrix has a match value of 10 for each identical nucleotide and -9 for each mismatch. The default gap creation penalty is -50 and the default gap extension penalty is -3 for each nucleotide.

10 The present invention also encompasses nucleotide sequences that are capable of hybridising selectively to the sequences presented herein, or any variant, fragment or derivative thereof, or to the complement of any of the above. Nucleotide sequences are preferably at least 15 nucleotides in length, more preferably at least 20, 30, 40 or 50 nucleotides in length.

15 HYBRIDISATION

The term "hybridization" as used herein shall include "the process by which a strand of nucleic acid joins with a complementary strand through base pairing" as well as the
20 process of amplification as carried out in polymerase chain reaction (PCR) technologies.

Nucleotide sequences of the invention capable of selectively hybridising to the nucleotide sequences presented herein, or to their complement, will be generally at least 75%,
25 preferably at least 85 or 90% and more preferably at least 95% or 98% homologous to the corresponding nucleotide sequences presented herein over a region of at least 20, preferably at least 25 or 30, for instance at least 40, 60 or 100 or more contiguous nucleotides. Preferred nucleotide sequences of the invention will comprise regions homologous to the nucleotide sequence set out in SEQ ID No 5 or SEQ ID No 7 or SEQ
30 ID No 8 of the sequence listings of the present invention preferably at least 80 or 90% and more preferably at least 95% homologous to the nucleotide sequence set out in SEQ ID No 5 or SEQ ID No 7 or SEQ ID No 8 of the sequence listings of the present invention.

The term "selectively hybridizable" means that the nucleotide sequence used as a probe is used under conditions where a target nucleotide sequence of the invention is found to hybridize to the probe at a level significantly above background. The background hybridization may occur because of other nucleotide sequences present, for example, in the cDNA or genomic DNA library being screened. In this event, background implies a level of signal generated by interaction between the probe and a non-specific DNA member of the library which is less than 10 fold, preferably less than 100 fold as intense as the specific interaction observed with the target DNA. The intensity of interaction may be measured, for example, by radiolabelling the probe, e.g. with ^{32}P .

Hybridization conditions are based on the melting temperature (T_m) of the nucleic acid binding complex, as taught in Berger and Kimmel (1987, Guide to Molecular Cloning Techniques, Methods in Enzymology, Vol 152, Academic Press, San Diego CA), and confer a defined "stringency" as explained below.

Maximum stringency typically occurs at about $T_m - 5^\circ\text{C}$ (5°C below the T_m of the probe); high stringency at about 5°C to 10°C below T_m ; intermediate stringency at about 10°C to 20°C below T_m ; and low stringency at about 20°C to 25°C below T_m . As will be understood by those of skill in the art, a maximum stringency hybridization can be used to identify or detect identical nucleotide sequences while an intermediate (or low) stringency hybridization can be used to identify or detect similar or related polynucleotide sequences.

In a preferred aspect, the present invention covers nucleotide sequences that can hybridise to the nucleotide sequence of the present invention under stringent conditions (e.g. 65°C and $0.1\times\text{SSC}$ { $1\times\text{SSC} = 0.15\text{ M NaCl}, 0.015\text{ M Na}_3\text{ Citrate pH } 7.0$). Where the nucleotide sequence of the invention is double-stranded, both strands of the duplex, either individually or in combination, are encompassed by the present invention. Where the nucleotide sequence is single-stranded, it is to be understood that the complementary sequence of that nucleotide sequence is also included within the scope of the present invention.

Nucleotide sequences which are not 100% homologous to the sequences of the present invention but fall within the scope of the invention can be obtained in a number of ways. Other variants of the sequences described herein may be obtained for example by probing DNA libraries made from a range of sources. In addition, other viral/bacterial, or cellular homologues particularly cellular homologues found in mammalian cells (e.g. rat, mouse, bovine and primate cells), may be obtained and such homologues and fragments thereof in general will be capable of selectively hybridising to the sequences shown in the sequence listing herein. Such sequences may be obtained by probing cDNA libraries made from or genomic DNA libraries from other animal species, and probing such libraries with probes comprising all or part of the nucleotide sequence set out in SEQ ID No 5 or SEQ ID No 7 or SEQ ID No 8 of the sequence listings of the present invention under conditions of medium to high stringency. Similar considerations apply to obtaining species homologues and allelic variants of the amino acid and/or nucleotide sequences of the present invention.

15 Variants and strain/species homologues may also be obtained using degenerate PCR which will use primers designed to target sequences within the variants and homologues encoding conserved amino acid sequences within the sequences of the present invention. Conserved sequences can be predicted, for example, by aligning the amino acid sequences from several variants/homologues. Sequence alignments can be performed using computer software known in the art. For example the GCG Wisconsin PileUp program is widely used. The primers used in degenerate PCR will contain one or more degenerate positions and will be used at stringency conditions lower than those used for cloning sequences with single sequence primers against known sequences.

25 Alternatively, such nucleotide sequences may be obtained by site directed mutagenesis of characterised sequences, such as the nucleotide sequence set out in SEQ ID No 5 or SEQ ID No 7 or SEQ ID No 8 of the sequence listings of the present invention. This may be useful where for example silent codon changes are required to sequences to optimise codon preferences for a particular host cell in which the nucleotide sequences are being expressed. Other sequence changes may be desired in order to introduce restriction enzyme recognition sites, or to alter the binding specificity of the ScFv Ab encoded by the nucleotide sequences.

The nucleotide sequences of the present invention may be used to produce a primer, e.g. a PCR primer, a primer for an alternative amplification reaction, a probe e.g. labelled with a revealing label by conventional means using radioactive or non-radioactive labels, or the nucleotide sequences may be cloned into vectors. Such primers, probes and other
5 fragments will be at least 15, preferably at least 20, for example at least 25, 30 or 40 nucleotides in length, and are also encompassed by the term nucleotide sequence of the invention as used herein.

The nucleotide sequences such as a DNA polynucleotides and probes according to the
10 invention may be produced recombinantly, synthetically, or by any means available to those of skill in the art. They may also be cloned by standard techniques.

In general, primers will be produced by synthetic means, involving a step wise manufacture of the desired nucleic acid sequence one nucleotide at a time. Techniques for
15 accomplishing this using automated techniques are readily available in the art.

Longer nucleotide sequences will generally be produced using recombinant means, for example using a PCR (polymerase chain reaction) cloning techniques. This will involve making a pair of primers (e.g. of about 15 to 30 nucleotides) flanking a region of the
20 targeting sequence which it is desired to clone, bringing the primers into contact with mRNA or cDNA obtained from an animal or human cell, performing a polymerase chain reaction (PCR) under conditions which bring about amplification of the desired region, isolating the amplified fragment (e.g. by purifying the reaction mixture on an agarose gel) and recovering the amplified DNA. The primers may be designed to contain suitable
25 restriction enzyme recognition sites so that the amplified DNA can be cloned into a suitable cloning vector

Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence, may be
30 used to clone and express the ScFv Ab. As will be understood by those of skill in the art, it may be advantageous to produce the ScFv Ab - encoding nucleotide sequences possessing non-naturally occurring codons. Codons preferred by a particular

prokaryotic or eukaryotic host (Murray E *et al* (1989) Nuc Acids Res 17:477-508) can be selected, for example, to increase the rate of the ScFv Ab expression or to produce recombinant RNA transcripts having desirable properties, such as a longer half-life, than transcripts produced from naturally occurring sequence.

5

In one embodiment of the present invention, the ScFv Ab is a recombinant ScFv Ab.

Preferably the recombinant ScFv Ab is prepared using a genetic vector.

10 VECTOR

As it is well known in the art, a vector is a tool that allows or facilitates the transfer of an entity from one environment to another. In accordance with the present invention, and by way of example, some vectors used in recombinant DNA techniques allow entities, such as a segment of DNA (such as a heterologous DNA segment, such as a heterologous cDNA segment), to be transferred into a host and/or a target cell for the purpose of replicating the vectors comprising the nucleotide sequences of the present invention and/or expressing the proteins of the invention encoded by the nucleotide sequences of the present invention. Examples of vectors used in recombinant DNA techniques include but are not limited to plasmids, chromosomes, artificial chromosomes or viruses.

20

The term "vector" includes expression vectors and/or transformation vectors.

25 The term "expression vector" means a construct capable of *in vivo* or *in vitro/ex vivo* expression.

The term "transformation vector" means a construct capable of being transferred from one species to another.

30

“NAKED DNA”

The vectors comprising nucleotide sequences encoding ScFv Abs of the present invention for use in affecting viral infections may be administered directly as “a naked
5 nucleic acid construct”, preferably further comprising flanking sequences homologous to the host cell genome.

As used herein, the term “naked DNA” refers to a plasmid comprising a nucleotide sequences encoding a ScFv Ab of the present invention together with a short promoter
10 region to control its production. It is called “naked” DNA because the plasmids are not carried in any delivery vehicle. When such a DNA plasmid enters a host cell, such as a eukaryotic cell, the proteins it encodes (such as the ScFv Ab) are transcribed and translated within the cell.

15 NON-VIRAL DELIVERY

Alternatively, the vectors comprising nucleotide sequences of the present invention may be introduced into suitable host cells using a variety of non-viral techniques known in the art, such as transfection, transformation, electroporation and biolistic
20 transformation.

As used herein, the term “transfection” refers to a process using a non-viral vector to deliver a gene to a target mammalian cell.

25 Typical transfection methods include electroporation, DNA biolistics, lipid-mediated transfection, compacted DNA-mediated transfection, liposomes, immunoliposomes, lipofectin, cationic agent-mediated, cationic facial amphiphiles (CFAs) (Nature Biotechnology 1996 14; 556), multivalent cations such as spermine, cationic lipids or polylysine, 1, 2,-bis (oleoyloxy)-3-(trimethylammonio) propane (DOTAP)-cholesterol
30 complexes (Wolff and Trubetskoy 1998 Nature Biotechnology 16: 421) and combinations thereof.

Uptake of naked nucleic acid constructs by mammalian cells is enhanced by several known transfection techniques for example those including the use of transfection agents. Example of these agents include cationic agents (for example calcium phosphate and DEAE-dextran) and lipofectants (for example lipofectamTM and transfectamTM). Typically, nucleic acid constructs are mixed with the transfection agent to produce a composition.

VIRAL VECTORS

Alternatively, the vectors comprising nucleotide sequences of the present invention may be introduced into suitable host cells using a variety of viral techniques which are known in the art, such as for example infection with recombinant viral vectors such as retroviruses, herpes simplex viruses and adenoviruses.

Preferably the vector is a recombinant viral vectors. Suitable recombinant viral vectors include but are not limited to adenovirus vectors, adeno-associated viral (AAV) vectors, herpes-virus vectors, a retroviral vector, lentiviral vectors, baculoviral vectors, pox viral vectors or parvovirus vectors (see Kestler et al 1999 Human Gene Ther 10(10):1619-32). In the case of viral vectors, gene delivery is mediated by viral infection of a target cell.

RETROVIRAL VECTORS

Examples of retroviruses include but are not limited to: murine leukemia virus (MLV), human immunodeficiency virus (HIV), equine infectious anaemia virus (EIAV), mouse mammary tumour virus (MMTV), Rous sarcoma virus (RSV), Fujinami sarcoma virus (FuSV), Moloney murine leukemia virus (Mo-MLV), FBR murine osteosarcoma virus (FBR MSV), Moloney murine sarcoma virus (Mo-MSV), Abelson murine leukemia virus (A-MLV), Avian myelocytomatosis virus-29 (MC29), and Avian erythroblastosis virus (AEV).

Preferred vectors for use in accordance with the present invention are recombinant viral vectors, in particular recombinant retroviral vectors (RRV) such as lentiviral vectors,.

The term "recombinant retroviral vector" (RRV) refers to a vector with sufficient retroviral genetic information to allow packaging of an RNA genome, in the presence of
5 packaging components, into a viral particle capable of infecting a target cell. Infection of the target cell includes reverse transcription and integration into the target cell genome. The RRV carries non-viral coding sequences which are to be delivered by the vector to the target cell. An RRV is incapable of independent replication to produce infectious retroviral particles within the final target cell. Usually the RRV lacks a
10 functional *gag-pol* and/or *env* gene and/or other genes essential for replication. The vector of the present invention may be configured as a split-intron vector. A split intron vector is described in PCT patent application WO 99/15683.

A detailed list of retroviruses may be found in Coffin *et al* ("Retroviruses" 1997 Cold
15 Spring Harbour Laboratory Press Eds: JM Coffin, SM Hughes, HE Varmus pp 758-763).

LENTIVIRAL VECTORS

20 Lentiviruses can be divided into primate and non-primate groups. Examples of primate lentiviruses include but are not limited to: the human immunodeficiency virus (HIV), the causative agent of human auto-immunodeficiency syndrome (AIDS), and the simian immunodeficiency virus (SIV). The non-primate lentiviral group includes the prototype
25 "slow virus" visna/maedi virus (VMV), as well as the related caprine arthritis-encephalitis virus (CAEV), equine infectious anaemia virus (EIAV) and the more recently described feline immunodeficiency virus (FIV) and bovine immunodeficiency virus (BIV).

A distinction between the lentivirus family and other types of retroviruses is that
30 lentiviruses have the capability to infect both dividing and non-dividing cells (Lewis *et al* 1992 EMBO. J 11: 3053-3058; Lewis and Emerman 1994 J. Virol. 68: 510-516). In

contrast, other retroviruses - such as MLV - are unable to infect non-dividing cells such as those that make up, for example, muscle, brain, lung and liver tissue.

ADENOVIRUSES

5

In one embodiment of the present invention, the features of adenoviruses may be combined with the genetic stability of retroviruses/lentiviruses which can be used to transduce target cells to become transient retroviral producer cells capable of stably infect neighbouring cells. Such retroviral producer cells which are engineered to
10 express a ScFv Ab of the present invention can be implanted in organisms such as animals or humans for use in the treatment of disease such as cancer.

POX VIRUSES

15 Preferred vectors for use in accordance with the present invention are recombinant pox viral vectors such as fowl pox virus (FPV), entomopox virus, vaccinia virus such as NYVAC, canarypox virus, MVA or other non-replicating viral vector systems such as those described for example in WO 95/30018.

20 HYBRID VIRAL VECTORS

In a further broad aspect, the present invention provides a hybrid viral vector system for *in vivo* delivery of a nucleotide sequence encoding a ScFc Ab of the present invention, which system comprises one or more primary viral vectors which encode a secondary
25 viral vector, the primary vector or vectors capable of infecting a first target cell and of expressing therein the secondary viral vector, which secondary vector is capable of transducing a secondary target cell.

Preferably the primary vector is obtainable from or is based on an adenoviral vector
30 and/or the secondary viral vector is obtainable from or is based on a retroviral vector preferably a lentiviral vector.

TARGETED VECTOR

The term "targeted vector" refers to a vector whose ability to infect/transfect/transduce a cell or to be expressed in a host and/or target cell is restricted to certain cell types
5 within the host organism, usually cells having a common or similar phenotype.

REPLICATION VECTORS

The nucleotide sequences encoding the ScFv Ab of the present invention may be
10 incorporated into a recombinant replicable vector. The vector may be used to replicate the nucleotide sequence in a compatible host cell. Thus in one embodiment of the present invention, the invention provides a method of making the ScFv Ab of the present invention by introducing a nucleotide sequence of the present invention into a replicable vector, introducing the vector into a compatible host cell, and growing the
15 host cell under conditions which bring about replication of the vector. The vector may be recovered from the host cell.

EXPRESSION VECTOR

20 Preferably, a nucleotide sequence of present invention which is inserted into a vector is operably linked to a control sequence that is capable of providing for the expression of the coding sequence, such as the coding sequence of the ScFv Ab of the present invention by the host cell, i.e. the vector is an expression vector. The ScFv Ab produced by a host recombinant cell may be secreted or may be contained intracellularly
25 depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing the ScFv Ab coding sequences can be designed with signal sequences which direct secretion of the ScFv Ab coding sequences through a particular prokaryotic or eukaryotic cell membrane.

EXPRESSION IN VITRO

The vectors of the present invention may be transformed or transfected into a suitable host cell and/or a target cell as described below to provide for expression of an ScFv Ab of the present invention. This process may comprise culturing a host cell and/or target cell transformed with an expression vector under conditions to provide for expression by the vector of a coding sequence encoding the ScFv Ab and optionally recovering the expressed ScFv Ab. The vectors may be for example, plasmid or virus vectors provided with an origin of replication, optionally a promoter for the expression of the said polynucleotide and optionally a regulator of the promoter. The vectors may contain one or more selectable marker genes, for example an ampicillin resistance gene in the case of a bacterial plasmid or a neomycin resistance gene for a mammalian vector. The expression of the ScFv Abs of the invention may be constitutive such that they are continually produced, or inducible, requiring a stimulus to initiate expression. In the case of inducible expression, ScFv Ab production can be initiated when required by, for example, addition of an inducer substance to the culture medium, for example dexamethasone or IPTG.

ScFv Ab CONSTRUCTS

FUSION PROTEINS

The ScFv Ab of the invention may also be produced as fusion proteins, for example to aid in extraction and purification. Examples of fusion protein partners include glutathione-S-transferase (GST), 6xHis, GAL4 (DNA binding and/or transcriptional activation domains) and β -galactosidase. Other examples of fusion protein partners include but are not limited to a fused recombinant ScFv Ab protein comprising an antigenic co-protein such as GST, β -galactosidase or the lipoprotein D from *Haemophilus influenzae* which are relatively large co-proteins, which solubilise and facilitate production and purification thereof. Alternatively, the fused protein may comprise a carrier protein such as bovine serum albumin (BSA) or keyhole limpet haemocyanin (KLH). In certain embodiments of the present invention, the marker

sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen Inc) and described in Gentz *et al* (1989 PNAS 86: 821-824). Such fusion proteins are readily expressable in yeast culture (as described in Mitchell *et al* 1993 Yeast 5: 715-723) and are easily purified by affinity chromatography.

5

Other recombinant constructions may join the ScFv Ab coding sequence to nucleotide sequence encoding a polypeptide domain which will facilitate purification of soluble proteins (Kroll *DJ et al* (1993) DNA Cell Biol 12:441-53). Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals (Porath J 10 (1992) Protein Expr Purif 3 -.26328 1), protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp, Seattle, WA).

15 It may also be convenient to include a proteolytic cleavage site between the fusion protein partner and the protein sequence of interest to allow removal of fusion protein sequences. By way of example, a fusion protein may also be engineered to contain a cleavage site located between the nucleotide sequence encoding the ScFv Ab and the heterologous protein sequence, so that the ScFv Ab may be cleaved and purified away 20 from the heterologous moiety. The inclusion of a cleavable linker sequence such as Factor XA or enterokinase (Invitrogen, San Diego, CA) between the purification domain and the ScFv Ab may also be useful to facilitate purification. Preferably the fusion protein will not hinder the binding specificity of the ScFv Ab comprising the amino acid sequence of the present invention.

25

In one preferred embodiment, the fusion protein comprises or encodes a secreted co-stimulatory molecule (SCM).

SCM FUSION PROTEINS

30

The secreted co-stimulatory molecule (SCM) of the invention may be an engineered fusion protein comprising a signal peptide for secretion from mammalian cells and at

least one further domain which acts as a co-stimulatory signal to a cell of the immune system. The use of combinations of SCMs containing different co-stimulatory domains may also envisaged. The ScFv Abs comprising the SCMs may be produced by expression of SCM-encoding genes in the autologous cells of the individual to be treated and hence any post-translational modifications added to the protein by the host cell are authentic and provide fully functional protein and appropriate pharmacokinetics.

WO-A-92/00092 describes truncated forms of B7-1, derived by placing a translation stop codon before the transmembrane domain, secreted from mammalian cells. In that particular case, a heterologous signal peptide from the Oncostatin M gene was used. WO-A92/00092 also describes fusion proteins which contain the extracellular domain of B7-1 fused to the Fc region of an immunoglobulin. Such molecules can bind to CD28 on T-cells and serve to stimulate T-cell proliferation. However such stimulation occurs only to a moderate extent unless the B7 or B7-derivative is immobilised on a solid surface.

Gerstmayer *et al.* (1997 J. Immunol. 158: 4584-4590) describes a fusion of B7-2 to an ScFv specific for ErbB2 followed by a myc epitope tag and polyhistidine tag which is secreted when expressed in the yeast *Pichia pastoris*. This molecule retained binding for antigen and co-stimulated proliferation of T-cells prestimulated with PMA and IL-2. However, glycosylation of such a molecule is of the yeast type, which is likely to lead to inappropriate pharmacokinetics in humans.

In accordance with the present invention, any suitable co-stimulatory domain(s) may be used. By way of example, co-stimulatory domains can be chosen from extracellular portions of the B7 family of cell-surface glycoproteins, including B7-1, B7-2 and B7-3 or other co-stimulatory cell surface glycoproteins such as but not limited to co-stimulatory receptor-ligand molecules including CD2/LFA-3, LFA-1/ICAM-1 and ICAM-3. Studies have demonstrated that T cell co-stimulation by monocytes is dependent on each of two receptor ligand pathways CD2/LFA-3 and LFA-1/ICAM-1 (Van Severter *et al* 1991 Eur J Immunol 21: 1711-1718). In addition, it has been

shown that ICAM-3, the third LFA-1 counterreceptor, is a co-stimulatory molecule for resting and activated T lymphocytes (Hernandez-Caselles *et al* 1993 Eur J Immunol 23: 2799-2806).

5 Other possible co-stimulatory molecules may include a novel glycoprotein receptor designated SLAM, has been identified which, when engaged, potentiates T-cell expansion in a CD28-independent manner and induces a Th0/Th1 cytokine production profile (Cocks *et al* 1995 Nature 376: 260-263).

10 CD6, a cell surface glycoprotein, has also been shown to function as a co-stimulatory and adhesion receptor on T cells. Four CD6 isoforms (CD6a, b, c, d) have been described (Kobarg *et al* 1997 Eur J Immunol 27: 2971-2980). A role for the very late antigen (VLA-4) integrin in the activation of human memory B cells has also been suggested (Silvy *et al* 1997 Eur J Immunol 27: 2757-2764). Endothelial cells also
 15 provide unique co-stimulatory signals that affect the phenotype of activated CD4+ T cells (Karmann *et al* 1996 Eur J Immunol 26: 610-617). A B3 protein, present on the surface of lipopolysaccharide-activated B cells, which can provide co-stimulation to resting T cells leading to a predominant release of interleukin-4 (IL-4) and IL-5 and negligible amounts of IL-2 and interferon gamma has been described (Vinay *et al* 1995
 20 J Biol Chem 270: 23429-23436). The co-expression of a novel co-stimulatory T cell antigen (A6H) on T cells and tumour cells has suggested a possible function related to common properties of these cells (Labuda *et al* 1995 Int Immunol 7: 1425-1432).

In one preferred embodiment of the invention, the co-stimulatory domain is a portion of
 25 B7-1 or B7-2, more preferably the complete extracellular portion of B7-1 or B7-2.

In one preferred embodiment the ScFv Ab of the present invention is formed by expression of a novel gene encoding a fusion protein containing the DACSM binding domain or domains and the co-stimulatory domain or domains. In the context of the
 30 present invention, the co-stimulatory domain is fused to the ScFv. The domains can be placed in the order (N-terminus to C-terminus): antigen-binding domain followed by co-stimulatory domain; or co-stimulatory domain followed by antigen-binding domain.

Preferably, the co-stimulatory domain is placed at the N-terminus followed by the antigen-binding domain. A signal peptide is included at the N-terminus, and may be for example the natural signal peptide of the co-stimulatory extracellular domain. The different domains may be separated by additional sequences, which may result from the inclusion of convenient restriction-enzyme cleavage sites in the novel gene to facilitate its construction, or serve as a peptide spacer between the domains, or serve as a flexible peptide linker or provide another function. Preferably the domains are separated by a flexible linker.

Two or more different genes encoding different SCMs may be used to achieve improved co-stimulation, or both co-stimulation of naïve T-cells and induction of memory responses. For example a gene encoding an SCM containing the B7-1 extracellular domain may be administered with a gene encoding an SCM containing the B7-2 extracellular domain.

QUANTITATION OF SCFV ANTIBODY PRODUCTION

Although the presence/absence of marker gene expression may suggest that the nucleotide sequence and/or its ScFv Ab is also present, its presence and expression may be confirmed by routine means. For example, if the ScFv Ab encoding nucleotide sequence is inserted within a marker gene sequence, recombinant cells containing the ScFv Ab coding regions may be identified by the absence of the marker gene function. Alternatively, a marker gene may be placed in tandem with a ScFv Ab encoding nucleotide sequence under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the ScFv Ab as well.

Additional methods to quantitate the expression of a particular molecule include radiolabeling (Melby PC *et al* 1993 J Immunol Methods 159:235-44) or biotinylating (Duplaa C *et al* 1993 Anal Biochem 229-36) nucleotides, coamplification of a control nucleic acid, and standard curves onto which the experimental results are interpolated. Quantitation of multiple samples may be speeded up by running the assay in an ELISA

format where the ScFv Ab of interest is presented in various dilutions and a spectrophotometric or calorimetric response gives rapid quantitation.

HOST/TARGET CELLS

5

Host and/or target cells comprising nucleotide sequences of the present invention may be used to express the ScFv Abs of the present invention under *in vitro*, *in vivo* and *ex vivo* conditions.

10

The term "host cell and/or target cell" includes any cell derivable from a suitable organism which a vector is capable of transfecting or transducing. Examples of host and/or target cells can include but are not limited to cells capable of expressing the ScFv Ab of the present invention under *in vitro*, *in vivo* and *ex vivo* conditions. Examples of such cells include but are not limited to macrophages, endothelial cells or combinations thereof. Further examples include respiratory airway epithelial cells, hepatocytes, muscle cells, cardiac myocytes, synoviocytes, primary mammary epithelial

15

cells and post-mitotically terminally differentiated non-replicating cells such as macrophages and/or neurons.

20

In a preferred embodiment, the cell is a mammalian cell.

In a highly preferred embodiment, the cell is a human cell.

25

The term "organism" includes any suitable organism. In a preferred embodiment, the organism is a mammal. In a highly preferred embodiment, the organism is a human.

30

Although the ScFv Ab of the invention may be produced using prokaryotic cells as host cells, it is preferred to use eukaryotic cells, for example yeast, insect or mammalian cells, in particular mammalian cells. Suitable host cells include bacteria such as *E. coli*, yeast, mammalian cell lines and other eukaryotic cell lines, for example insect Sf9 cells.

The present invention also provides a method comprising transforming a host and/or target cell with a or the nucleotide sequence(s) of the present invention.

5 The term "transformed cell" means a host cell and/or a target cell having a modified genetic structure. With the present invention, a cell has a modified genetic structure when a vector according to the present invention has been introduced into the cell.

Host cells and/or a target cells may be cultured under suitable conditions which allow expression of the ScFv Ab of the invention.

10

The present invention also provides a method comprising culturing a transformed host cell - which cell has been transformed with a or the nucleotide sequence(s) according to the present invention under conditions suitable for the expression of the ScFv Ab encoded by said nucleotide sequence(s).

15

The present invention also provides a method comprising culturing a transformed host cell - which cell has been transformed with a or the nucleotide sequence(s) according to the present invention or a derivative, homologue, variant or fragment thereof - under conditions suitable for the expression of the ScFv Ab encoded by said nucleotide sequence(s); and then recovering said ScFv Ab from the transformed host cell culture.

20

The ScFv Ab of the present invention can be extracted from host cells by a variety of techniques known in the art, including enzymatic, chemical and/or osmotic lysis and physical disruption. The ScFv Ab may be purified and isolated in a manner known *per se*.

25

REGULATION OF EXPRESSION IN VITRO/ VIVO/EX VIVO

30

The present invention also encompasses gene therapy whereby the ScFv Ab encoding nucleotide sequence(s) of the present invention is regulated *in vitro/in vivo/ex vivo*. For example, expression regulation may be accomplished by administering compounds that bind to the ScFv Ab encoding nucleotide sequence(s) of the present invention, or control regions associated with the ScFv Ab encoding nucleotide sequence of the

present invention, or its corresponding RNA transcript to modify the rate of transcription or translation.

CONTROL SEQUENCES

5

Control sequences operably linked to sequences encoding the ScFv Ab of the present invention include promoters/enhancers and other expression regulation signals. These control sequences may be selected to be compatible with the host cell and/or target cell in which the expression vector is designed to be used. The control sequences may be modified, for example by the addition of further transcriptional regulatory elements to make the level of transcription directed by the control sequences more responsive to transcriptional modulators.

10

OPERABLY LINKED

15

The term "operably linked" means that the components described are in a relationship permitting them to function in their intended manner. A regulatory sequence "operably linked" to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under condition compatible with the control sequences.

20

Preferably the nucleotide sequence of the present invention is operably linked to a transcription unit.

25

The term "transcription unit(s)" as described herein are regions of nucleic acid containing coding sequences and the signals for achieving expression of those coding sequences independently of any other coding sequences. Thus, each transcription unit generally comprises at least a promoter, an optional enhancer and a polyadenylation signal.

30

PROMOTERS

The term promoter is well-known in the art and is used in the normal sense of the art, e.g. as an RNA polymerase binding site. The term encompasses nucleic acid regions
5 ranging in size and complexity from minimal promoters to promoters including upstream elements and enhancers.

The promoter is typically selected from promoters which are functional in mammalian, cells, although prokaryotic promoters and promoters functional in other eukaryotic cells
10 may be used. The promoter is typically derived from promoter sequences of viral or eukaryotic genes. For example, it may be a promoter derived from the genome of a cell in which expression is to occur. With respect to eukaryotic promoters, they may be promoters that function in a ubiquitous manner (such as promoters of α -actin, β -actin, tubulin) or, alternatively, a tissue-specific manner (such as promoters of the genes for
15 pyruvate kinase).

HYPOXIC PROMOTERS/ENHANCERS

The enhancer and/or promoter may be preferentially active in a hypoxic or ischaemic or
20 low glucose environment, such that the ScFv Ab encoding nucleotide sequence(s) is preferentially expressed in the particular tissues of interest, such as in the environment of a tumour, arthritic joint or other sites of ischaemia. Thus, any significant biological effect or deleterious effect of the ScFv Ab encoding nucleotide sequence(s) on the individual being treated may be reduced or eliminated. The enhancer element or other
25 elements conferring regulated expression may be present in multiple copies. Likewise, or in addition, the enhancer and/or promoter may be preferentially active in one or more specific cell types - such as any one or more of macrophages, endothelial cells or combinations thereof. Further examples may include but are not limited to respiratory airway epithelial cells, hepatocytes, muscle cells, cardiac myocytes, synoviocytes,
30 primary mammary epithelial cells and post-mitotically terminally differentiated non-replicating cells such as macrophages and/or neurons.

TISSUE-SPECIFIC PROMOTERS

The promoters of the present invention may be tissue-specific promoters. Examples of suitable tissue restricted promoters/enhancers are those which are highly active in tumour cells such as a promoter/enhancer from a *MUC1* gene, a *CEA* gene or a *5T4* antigen gene. Examples of temporally restricted promoters/enhancers are those which are responsive to ischaemia and/or hypoxia, such as hypoxia response elements or the promoter/enhancer of a *grp78* or a *grp94* gene. The alpha fetoprotein (AFP) promoter is also a tumour-specific promoter. One preferred promoter-enhancer combination is a human cytomegalovirus (hCMV) major immediate early (MIE) promoter/enhancer combination.

Preferably the promoters of the present invention are tissue specific. That is, they are capable of driving transcription of a ScFv Ab encoding nucleotide sequence(s) in one tissue while remaining largely "silent" in other tissue types.

The term "tissue specific" means a promoter which is not restricted in activity to a single tissue type but which nevertheless shows selectivity in that they may be active in one group of tissues and less active or silent in another group. A desirable characteristic of the promoters of the present invention is that they possess a relatively low activity in the absence of activated hypoxia-regulated enhancer elements, even in the target tissue. One means of achieving this is to use "silencer" elements which suppress the activity of a selected promoter in the absence of hypoxia.

The term "hypoxia" means a condition under which a particular organ or tissue receives an inadequate supply of oxygen.

The level of expression of a or the ScFv Ab encoding nucleotide sequence(s) under the control of a particular promoter may be modulated by manipulating the promoter region. For example, different domains within a promoter region may possess different gene regulatory activities. The roles of these different regions are typically assessed using vector constructs having different variants of the promoter with specific regions

deleted (that is, deletion analysis). This approach may be used to identify, for example, the smallest region capable of conferring tissue specificity or the smallest region conferring hypoxia sensitivity.

5 A number of tissue specific promoters, described above, may be particularly advantageous in practising the present invention. In most instances, these promoters may be isolated as convenient restriction digestion fragments suitable for cloning in a selected vector. Alternatively, promoter fragments may be isolated using the polymerase chain reaction. Cloning of the amplified fragments may be facilitated by
10 incorporating restriction sites at the 5' end of the primers.

INDUCIBLE PROMOTERS

The promoters of the present invention may also be promoters that respond to specific
15 stimuli, for example promoters that bind steroid hormone receptors. Viral promoters may also be used, for example the Moloney murine leukaemia virus long terminal repeat (MMLV LTR) promoter, the rous sarcoma virus (RSV) LTR promoter or the human cytomegalovirus (CMV) IE promoter.

20 It may also be advantageous for the promoters to be inducible so that the levels of expression of the heterologous gene can be regulated during the life-time of the cell. Inducible means that the levels of expression obtained using the promoter can be regulated.

25 ENHANCER

In addition, any of these promoters may be modified by the addition of further regulatory sequences, for example enhancer sequences.. Chimeric promoters may also be used comprising sequence elements from two or more different promoters described
30 above.

The term "enhancer" includes a DNA sequence which binds to other protein components of the transcription initiation complex and thus facilitates the initiation of transcription directed by its associated promoter.

- 5 The *in vitro/ in vivo/ ex vivo* expression of the ScFv Ab of the present invention may be used in combination with a protein of interest (POI) or a nucleotide sequence of interest (NOI) encoding same.

COMBINATION WITH POIs/NOIs

- 10 The ScFv Ab of the present invention or nucleotide sequence encoding same may be used in combination with a POI, such as a pro-drug activating enzyme either directly or by vector delivery to, for example, a target cell or target tissue. Instead of or as well as being selectively expressed in target tissues, the ScFv Ab of the present invention or
15 nucleotide sequence encoding same may be used in combination with another POI such as a pro-drug activation enzyme or enzymes or with a nucleotide sequences of interest (NOI) or NOIs which encode a pro-drug activation enzyme or enzymes. These pro-drug activation enzyme or enzymes may have no significant effect or no deleterious effect until the individual is treated with one or more pro-drugs upon which the
20 appropriate pro-drug enzyme or enzymes act. In the presence of the active POI or NOI encoding same, treatment of an individual with the appropriate pro-drug may lead to enhanced reduction in the disease condition such as a reduction in tumour growth or survival.

25 PRO-DRUG POIs

- A POI, such as a pro-drug activating enzyme, may be delivered to a disease site, such as a tumour site for the treatment of a cancer. In each case, a suitable pro-drug is used in the treatment of the patient in combination with the appropriate pro-drug activating
30 enzyme. An appropriate pro-drug may be administered in conjunction with the ScFv Ab or vector comprising the nucleotide sequence encoding same. Examples of pro-drugs include: etoposide phosphate (with alkaline phosphatase, Senter *et al* 1988 Proc

Natl Acad Sci 85: 4842-4846); 5-fluorocytosine (with cytosine deaminase, Mullen *et al* 1994 Cancer Res 54: 1503-1506); Doxorubicin-N-p-hydroxyphenoxycetamide (with Penicillin-V-Amidase, Kerr *et al* 1990 Cancer Immunol Immunother 31: 202-206); Para-N-bis(2-chloroethyl) aminobenzoyl glutamate (with carboxypeptidase G2);
 5 Cephalosporin nitrogen mustard carbamates (with β -lactamase); SR4233 (with P450 Reductase); Ganciclovir (with HSV thymidine kinase, Borrelli *et al* 1988 Proc Natl Acad Sci 85: 7572-7576); mustard pro-drugs with nitroreductase (Friedlos *et al* 1997 J Med Chem 40: 1270-1275) and Cyclophosphamide (with P450 Chen *et al* 1996 Cancer Res 56: 1331-1340).

10

Examples of suitable pro-drug activation enzymes for use in the invention include a thymidine phosphorylase which activates the 5-fluoro-uracil pro-drugs capecitabine and fluturone; thymidine kinase from Herpes Simplex Virus which activates ganciclovir; a cytochrome P450 which activates a pro-drug such as cyclophosphamide to a DNA
 15 damaging agent; and cytosine deaminase which activates 5-fluorocytosine. Preferably, a pro-drug activating enzyme of human origin is used.

POIs AND NOIs

20 Other suitable proteins of interest (POIs) or NOIs encoding same for use in the present invention include those that are of therapeutic and/or diagnostic application such as, but are not limited to: sequences encoding cytokines, chemokines, hormones, antibodies, engineered immunoglobulin-like molecules, a single chain antibody, fusion proteins, enzymes, immune co-stimulatory molecules, immunomodulatory molecules, anti-sense
 25 RNA, a transdominant negative mutant of a target protein, a toxin, a conditional toxin, an antigen, a tumour suppressor protein and growth factors, membrane proteins, vasoactive proteins and peptides, anti-viral proteins and ribozymes, and derivatives thereof (such as with an associated reporter group). When included, the POIs or NOIs encoding same may be typically operatively linked to a suitable promoter, which may
 30 be a promoter driving expression of a ribozyme(s), or a different promoter or promoters, such as in one or more specific cell types.

BYSTANDER EFFECT

The POI and/or NOI encoding same may be proteins which are secreted from a cell. Alternatively the POI expression products are not secreted and are active within the cell. In either event, it is preferred for the POI expression product to demonstrate a bystander effector or a distant bystander effect; that is the production of the expression product in one cell leading to the killing of additional, related cells, either neighbouring or distant (e.g. metastatic), which possess a common phenotype.

Suitable POIs or NOIs encoding same for use in the present invention in the treatment or prophylaxis of cancer include proteins which: destroy the target cell (for example a ribosomal toxin), act as: tumour suppressors (such as wild-type p53); activators of anti-tumour immune mechanisms (such as cytokines, co-stimulatory molecules and immunoglobulins); inhibitors of angiogenesis; or which provide enhanced drug sensitivity (such as pro-drug activation enzymes); indirectly stimulate destruction of target cell by natural effector cells (for example, strong antigen to stimulate the immune system or convert a precursor substance to a toxic substance which destroys the target cell (for example a prodrug activating enzyme). Encoded proteins could also destroy bystander tumour cells (for example with secreted antitumour antibody-ribosomal toxin fusion protein), indirectly stimulate destruction of bystander tumour cells (for example cytokines to stimulate the immune system or procoagulant proteins causing local vascular occlusion) or convert a precursor substance to a toxic substance which destroys bystander tumour cells (eg an enzyme which activates a prodrug to a diffusible drug).

Also, the delivery of NOI(s) encoding antisense transcripts or ribozymes which interfere with expression of cellular genes for tumour persistence (for example against aberrant *myc* transcripts in Burkitts lymphoma or against *bcr-abl* transcripts in chronic myeloid leukemia. The use of combinations of such POIs and/or NOIs encoding same is also envisaged.

30

Examples of hypoxia regulatable therapeutic NOIs can be found in PCT/GB95/00322 (WO-A-9521927).

SCFV Ab COUPLING

The ScFv Ab of the present invention can be coupled to other molecules using standard methods. The amino and carboxyl termini of ScFv Ab may be isotopically and nonisotopically labeled with many techniques, for example radiolabeling using conventional techniques (tyrosine residues- chloramine T, iodogen, lactoperoxidase; lysine residues- Bolton-Hunter reagent). These coupling techniques are well known to those skilled in the art. The coupling technique is chosen on the basis of the functional groups available on the amino acids including, but not limited to amino, sulfhydryl, carboxyl, amide, phenol, and imidazole. Various reagents used to effect these couplings include among others, glutaraldehyde, diazotized benzidine, carbodiimide, and p-benzoquinone.

CHEMICAL COUPLING

The ScFv Ab of the present invention may be chemically coupled to isotopes, enzymes, carrier proteins, cytotoxic agents, fluorescent molecules, radioactive nucleotides and other compounds for a variety of applications including but not limited to imaging/prognosis, diagnosis and/or therapy. The efficiency of the coupling reaction is determined using different techniques appropriate for the specific reaction. For example, radiolabeling of an ScFv Ab peptide with ^{125}I is accomplished using chloramine T and Na^{125}I of high specific activity. The reaction is terminated with sodium metabisulfite and the mixture is desalted on disposable columns. The labeled peptide is eluted from the column and fractions are collected. Aliquots are removed from each fraction and radioactivity measured in a gamma counter. In this manner, the unreacted Na^{125}I is separated from the labeled ScFv Ab. The peptide fractions with the highest specific radioactivity are stored for subsequent use such as analysis of the ability to bind to a ScFv Ab.

IMAGING

The use of labelled ScFv Abs of the present invention with short lived isotopes enables visualization quantitation of DACSM binding sites *in vivo* using autoradiographic, or
5 modern radiographic or other membrane binding techniques such as positron emission tomography in order to locate tumours with ScFv Ab binding sites. This application provides important diagnostic and/or prognostic research tools.

CONJUGATES

10

In other embodiments, the ScFv Ab of the invention is coupled to a scintigraphic radiolabel, a cytotoxic compound or radioisotope, an enzyme for converting a non-toxic prodrug into a cytotoxic drug, a compound for activating the immune system in order to target the resulting conjugate to a disease site such as a colon tumour, or a cell-
15 stimulating compound. Such conjugates have a "binding portion", which consists of the ScFv Ab of the invention, and a "functional portion", which consists of the radiolabel, toxin or enzyme. Different ScFv Abs can be synthesized for use in several applications including but not limited to the linkage of a ScFv Ab to cytotoxic agents for targeted killing of cells that bind the ScFv Ab.

20

The ScFv Ab may alternatively be used alone in order simply to block the activity of the DACSM, particularly by physically interfering with its binding of another compound.

25

The binding portion and the functional portion of the conjugate (if also a peptide or polypeptide) may be linked together by any of the conventional ways of cross linking polypeptides, such as those generally described in O'Sullivan *et al* (Anal. Biochem 1979: 100, 100-108). For example, one portion may be enriched with thiol groups and the other portion reacted with a bifunctional agent capable of reacting with those thiol
30 groups, for example the N-hydroxysuccinimide ester of iodoacetic acid (NHIA) or N-succinimidyl-3-(2-pyridyldithio)propionate (SPDP). Amide and thioether bonds, for

example achieved with *m*-maleimidobenzoyl-N-hydroxysuccinimide ester, are generally more stable *in vivo* than disulphide bonds.

Alternatively, if the binding portion contains carbohydrates, such as would be the case for an antibody or some antibody fragments, the functional portion may be linked via the carbohydrate portion using the linking technology in EP 0 088 695.

The functional portion of the conjugate may be an enzyme for converting a non-toxic prodrug into a toxic drug, for example the conjugates of Bagshawe and his colleagues (Bagshawe (1987) Br. J. Cancer 56, 531; Bagshawe *et al* (Br. J. Cancer 1988: 58, 700); WO 88/07378) or cyanide-releasing systems (WO 91/11201).

It may not be necessary for the whole enzyme to be present in the conjugate but, of course, the catalytic portion must be present. So-called "abzymes" may be used, where a ScFv Ab is raised to a compound involved in the reaction one wishes to catalyse, usually the reactive intermediate state. The resulting antibody can then function as an enzyme for the reaction.

The conjugate may be purified by size exclusion or affinity chromatography, and tested for dual biological activities. The antigen immunoreactivity may be measured using an enzyme-linked immunosorbent assay (ELISA) with immobilised antigen and in a live cell radio-immunoassay. An enzyme assay may be used for β -glucosidase using a substrate which changes in absorbance when the glucose residues are hydrolysed, such as *o*NPG (*o*-nitrophenyl- β -D-glucopyranoside), liberating 2-nitrophenol which is measured spectrophotometrically at 405 nm.

The stability of the conjugate may be tested *in vitro* initially by incubating at 37°C in serum, followed by size exclusion FPLC analysis. Stability *in vivo* can be tested in the same way in mice by analysing the serum at various times after injection of the conjugate. In addition, it is possible to radiolabel the ScFv Ab with ^{125}I , and the enzyme with ^{131}I before conjugation, and to determine the biodistribution of the conjugate, free ScFv Ab and free enzyme in mice.

Alternatively, the conjugate may be produced as a fusion compound by recombinant DNA techniques whereby a length of DNA comprises respective regions encoding the two portions of the conjugate either adjacent to one another or separated by a region encoding a linker peptide which does not destroy the desired properties of the conjugate.

Conceivably, two of the functional portions of the compound may overlap wholly or partly. The DNA is then expressed in a suitable host in known ways.

10 DIAGNOSTIC KITS

The present invention also includes diagnostic methods and kits for detection and measurement of DACSM in biological fluids and tissues, and for localization of a DACSM in tissues. The ScFv Ab of the present invention that possess high binding specificity can be used to establish easy to use kits for rapid, reliable, sensitive, and specific measurement and localization of a DACSM in extracts of plasma, urine, tissues, and in cell culture media. The ScFv Ab of the present invention may also be used in a diagnostic method and kit to permit detection of circulating DACSMs which, in certain situations, may indicate the progression of a disease state such as the spread of micrometastases by primary tumours *in situ*.

These kits may include but are not limited to the following techniques; competitive and non-competitive assays, radioimmunoassay, bioluminescence and chemiluminescence assays, fluorometric assays, sandwich assays, immunoradiometric assays, dot blots, enzyme linked assays including ELISA, microtiter plates, antibody coated strips or dipsticks for rapid monitoring of urine or blood, and immunocytochemistry. For each kit the range, sensitivity, precision, reliability, specificity and reproducibility of the assay are established. Intraassay and interassay variation is established at 20%, 50% and 80% points on the standard curves of displacement or activity.

One example of an assay kit commonly used in research and in the clinic is a radioimmunoassay (RIA) kit. After successful radioiodination and purification of a

ScFv Ab, the antiserum possessing the highest titer is added at several dilutions to tubes containing a relatively constant amount of radioactivity, such as 10,000 cpm, in a suitable buffer system. Other tubes contain buffer or preimmune serum to determine the non-specific binding. After incubation at 4°C for 24 hours, protein A is added and the tubes are vortexed, incubated at room temperature for 90 minutes, and centrifuged at approximately 2000-2500 times g at 4°C to precipitate the complexes of antiserum bound to the labeled ScFv Ab. The supernatant is removed by aspiration and the radioactivity in the pellets counted in a gamma counter. The antiserum dilution that binds approximately 10 to 40 % of the labeled ScFv Ab after subtraction of the non-specific binding is further characterized.

IMMUNOHISTOCHEMISTRY

An immunohistochemistry kit may also be used for localization of DACSM in tissues and cells. This immunohistochemistry kit provides instructions, a ScFv Ab, and possibly blocking serum and secondary antiserum linked to a fluorescent molecule such as fluorescein isothiocyanate, or to some other reagent used to visualize the primary antiserum. Immunohistochemistry techniques are well known to those skilled in the art. This immunohistochemistry kit permits localization of a DACSM in tissue sections and cultured cells using both light and electron microscopy. It is used for both research and clinical purposes. For example, tumours are biopsied or collected and tissue sections cut with a microtome to examine sites of DACSM production. Such information is useful for diagnostic and possibly therapeutic purposes in the detection and treatment of diseases such as cancer.

COMBINATION THERAPY

The ScFv Abs of the present invention may be used in combination with other compositions and procedures for the treatment of diseases. By way of example, the ScFv Abs may also be used in combination with conventional treatments of diseases such as cancer. By ways of further example, a tumor may be treated conventionally with surgery, radiation or chemotherapy combined with a ScFv Ab or a ScFv Ab may

be subsequently administered to the patient to extend the dormancy of micrometastases and to stabilize any residual primary tumor.

ScFv Ab DELIVERY

5

The ScFv Ab can be delivered with a therapeutically effective agent at the same moment in time and at the same site. Alternatively, the ScFv Ab and the therapeutically effective agent may be delivered at a different time and to a different site. The ScFv Ab and the therapeutically effective agent may even be delivered in the same delivery
10 vehicle for the prevention and/or treatment of a disease condition such as cancer.

Therapeutic strategies based on the use of the ScFv Ab include the recruitment and activation of T cells by using a fusion of an DACSM reactive ScFv Ab fragment with the bacterial superantigen staphylococcal enterotoxin (Dohlsten *et al* 1994) or by using
15 bispecific antibodies, directed to both DACSM and the T-cell CD3 antigen (Kroesen *et al* 1994). Anti-DACSM antibodies may also be conjugated to different bacterial toxins to yield potent immunotoxins (LeMaistre *et al* 1987; Zimmermann *et al* 1997).

ScFv Abs may be used in combination with cytotoxic agents for the prevention and/or
20 treatment of disease states such as angiogenesis and/or cancer. Cytotoxic agents such as ricin, linked to ScFv Ab may provide a tool for the destruction of cells that bind the ScFv Ab. These cells may be found in many locations, including but not limited to, micrometastases and primary tumours.

25 SCREENS

The ScFv Ab of the present invention or a derivative or homologue thereof and/or a cell line that expresses the ScFv Ab of the present invention or a derivative or homologue thereof may be used to screen for agents (such as peptides, organic or inorganic
30 molecules) capable of affecting the binding specificity of the ScFv Ab.

In one embodiment, the screens of the present invention may identify agonists and/or antagonists of the ScFv Ab of the present invention.

5 In another embodiment, the ScFv Ab of the present invention may be used in a variety of drug screening techniques. The ScFv Ab employed in such a test may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The abolition of ScFv Ab binding specificity or the formation of binding complexes between the ScFv Ab and the agent being tested may be measured.

10 Another technique for screening provides for high throughput screening (HTS) of agents having suitable binding affinity to the ScFv Abs and is based upon the method described in detail in WO 84/03564.

15 It is expected that the assay methods of the present invention will be suitable for both small and large-scale screening of test compounds as well as in quantitative assays.

PHAGE DISPLAY SCREENS

20 Phage display may be employed in the identification of agents, such as a DACSM that is engageable by the ScFv Ab of the present invention. Phage display is a protocol of molecular screening which utilises recombinant bacteriophage. The technology involves transforming bacteriophage with a nucleotide sequence encoding an appropriate ligand (such as a candidate DACSM) which capable of reacting with a target ScFv Ab (or a derivative or homologue thereof) or the nucleotide sequence (or a
25 derivative or homologue thereof) encoding same. The transformed bacteriophage (which preferably is tethered to a solid support) expresses the appropriate ligand (such as the candidate agent) and displays it on their phage coat. The entity or entities (such as cells) bearing the target ScFv Ab molecules which recognises the candidate DACSM are isolated and amplified. The successful candidate DACSM is then characterised.

30

The targeting of cells expressing DACSM with a ScFv Ab of the present invention facilitates the development of ScFv Abs to modulate the activity of cells expressing the

DACSM

In another embodiment of the present invention, an ScFv Ab library may be used to screen for antibodies against specific DACSMs. By way of example, a bacteriophage
5 may be transformed with a nucleotide sequence encoding an appropriate ligand (such as a candidate ScFv Ab) which is capable of reacting with a target DACSM (or a derivative or homologue thereof) or the nucleotide sequence (or a derivative or homologue thereof) encoding same. The transformed bacteriophage (which preferably is tethered to a solid support) expresses the appropriate ligand (such as the candidate
10 ScFv Ab) and displays it on their phage coat. The entity or entities (such as cells) bearing the target DACSM molecules which recognises the candidate ScFv Ab are isolated and amplified. The successful candidate ScFv Ab is then characterised.

By way of further example, a human ScFv fragment library called "the Griffin-1
15 library" has been constructed by recloning synthetic heavy and light chain variable regions (VH and VL) from the lox library vectors into the phagemid vector pHEN2. Modifications in the elution and screening procedures an result in the successful screening of phage display libraries for ScFv antibodies against a large variety of DACSMs (see de Bruin et al 1999, Nature Biotechnology 17: 397-399).

20 Phage display has advantages over standard affinity ligand screening technologies. The phage surface displays the candidate agent in a three dimensional configuration, more closely resembling its naturally occuring conformation. This allows for more specific and higher affinity binding for screening purposes.

25

ASSAYS FOR MIMETICS

The positive identification of either a DACSM or a ScFv Ab using phage display may facilitate the use of combinatorial libraries to identify mimetics capable of acting in the
30 same or a similiar manner. Such mimetics can be administered alone or in combination with other therapeutics for the treatment of diseases associated with the DACSM of the present invention.

DOSAGE

The dosage of the ScFv Ab of the present invention will depend on the disease state or condition being treated and other clinical factors such as weight and condition of the human or animal and the route of administration of the compound. Depending upon the half-life of the ScFv Ab in the particular animal or human, the ScFv Ab can be administered between several times per day to once a week. It is to be understood that the present invention has application for both human and veterinary use. The methods of the present invention contemplate single as well as multiple administrations, given either simultaneously or over an extended period of time.

FORMULATIONS

Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampoules and vials, and may be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example, water for injections, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules and tablets of the kind previously described.

The ScFv Ab of the present invention may be effective in preventing and/or treating diseases such as cancer related diseases. The present invention includes the method of treating diseases such as cancer related disease with an effective amount of a ScFv Ab of the present invention. The ScFv Ab of the present invention can be provided as a synthetic peptide or an isolated and substantially purified proteins or protein fragments or a combination thereof in pharmaceutically acceptable compositions using formulation methods known to those of ordinary skill in the art. These compositions can be administered by standard routes. These include but are not limited to: oral,

rectal, ophthalmic (including intravitreal or intracameral), nasal, topical (including buccal and sublingual), intrauterine, vaginal or parenteral (including subcutaneous, intraperitoneal, intramuscular, intravenous, intradermal, intracranial, intratracheal, and epidural) transdermal, intraperitoneal, intracranial, intracerebroventricular, intracerebral, intravaginal, intrauterine, or parenteral (e.g., intravenous, intraspinal, subcutaneous or intramuscular) routes.

The ScFv Ab formulations may conveniently be presented in unit dosage form and may be prepared by conventional pharmaceutical techniques. Such techniques include the step of bringing into association the active ingredient and the pharmaceutical carrier(s) or excipient(s). In general, the formulations are prepared by uniformly and intimately bringing into association the active ingredient with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.

In addition, the ScFv Abs of the present invention may be incorporated into biodegradable polymers allowing for sustained release of the compound, the polymers being implanted in the vicinity of where drug delivery is desired, for example, at the site of a tumor or implanted so that the ScFv Ab is slowly released systemically. The biodegradable polymers and their use are described, for example, in detail in Brem *et al* (J. Neurosurg 1991 74:441-446). Osmotic minipumps may also be used to provide controlled delivery of high concentrations of ScFv Abs through cannulae to the site of interest, such as directly into a metastatic growth or into the vascular supply to that tumor.

The ScFv Abs of the present invention may be linked to cytotoxic agents which are infused in a manner designed to maximize delivery to the desired location. For example, ricin-linked high affinity ScFv Abs are delivered through a cannula into vessels supplying the target site or directly into the target. Such agents are also delivered in a controlled manner through osmotic pumps coupled to infusion cannulae.

Preferred unit dosage formulations are those containing a daily dose or unit, daily sub-dose, as herein above recited, or an appropriate fraction thereof, of the administered ingredient. It should be understood that in addition to the ingredients, particularly

mentioned above, the formulations of the present invention may include other agents conventional in the art having regard to the type of formulation in question.

5 The ScFv Ab conjugates may be administered in any suitable way, usually parenterally, for example intravenously or intraperitoneally, in standard sterile, non-pyrogenic formulations of diluents and carriers, for example isotonic saline (when administered intravenously). Once the ScFv Ab conjugate has bound to the target cells and been cleared from the bloodstream (if necessary), which typically takes a day or so, the pro-
10 drug is administered, usually as a single infused dose, or the tumour is imaged. If needed, because the ScFv Ab conjugate may be immunogenic, cyclosporin or some other immunosuppressant can be administered to provide a longer period for treatment but usually this will not be necessary.

15 The timing between administrations of the ScFv Ab conjugate and pro-drug may be optimised in a routine way since disease/normal tissue ratios of conjugate (at least following intravenous delivery) are highest after about 4-6 days, whereas at this time the absolute amount of conjugate bound to the DACSM, in terms of percent of injected dose per gram, is lower than at earlier times.

20 Therefore, the optimum interval between administration of the ScFv Ab conjugate and the pro-drug will be a compromise between peak concentration of the enzyme at the disease site and the best distribution ratio between disease and normal tissues. The dosage of the ScFv Ab conjugate will be chosen by the physician according to the usual criteria. At least in the case of methods employing a targeted enzyme such as β -
25 glucosidase and intravenous amygdalin as the toxic pro-drug, 1 to 50 daily doses of 0.1 to 10.0 grams per square metre of body surface area, preferably 1.0-5.0 g/m² are likely to be appropriate. For oral therapy, three doses per day of 0.05 to 10.0g, preferably 1.0-5.0g, for one to fifty days may be appropriate. The dosage of the ScFv Ab conjugate will similarly be chosen according to normal criteria, particularly with
30 reference to the type, stage and location of the disease tissue and the weight of the patient. The duration of treatment will depend in part upon the rapidity and extent of any immune reaction to the ScFv Ab conjugate.

The functional portion of the ScFv Ab conjugate, when the the ScFv Ab conjugate is used for diagnosis, usually comprises and may consist of a radioactive atom for scintigraphic studies, for example technetium 99m (^{99m}Tc) or iodine-123 (^{123}I), or a spin label for nuclear magnetic resonance (nmr) imaging (also known as magnetic resonance imaging, mri), such as iodine-123 again, iodine-313, indium-111, fluorine-19, carbon-13, nitrogen-15, oxygen-17, gadolinium, manganese or iron.

When used in a compound for selective destruction of, for example, the tumour, the functional portion of the ScFv Ab may comprise a highly radioactive atom, such as iodine-131, rhenium-186, rhenium-188, yttrium-90 or lead-212, which emits enough energy to destroy neighbouring cells, or a cytotoxic chemical compound such as methotrexate, adriamycin, vinca alkaloids (vincristine, vinblastine, etoposide), daunorubicin or other intercalating agents.

The radio- or other labels may be incorporated in the ScFv Ab conjugate in known ways. For example, the peptide may be biosynthesised or may be synthesised by chemical amino acid synthesis using suitable amino acid precursors involving, for example, fluorine-19 in place of hydrogen. Labels such as ^{99m}Tc , ^{123}I , ^{186}Rh , ^{188}Rh and ^{111}In can be attached via a cysteine residue in the peptide. Yttrium-90 can be attached via a lysine residue. The IODOGEN method (Fraker *et al* (1978) Biochem. Biophys. Res. Commun. 80: 49-57 can be used to incorporate iodine-123. "Monoclonal Antibodies in Immunoscintigraphy" (Chatal, CRC Press 1989) describes other methods in detail.

PHARMACEUTICAL COMPOSITIONS

In one aspect, the present invention provides a pharmaceutical composition, which comprises an ScFv Ab according to the present invention and optionally a pharmaceutically acceptable carrier, diluent or excipient (including combinations thereof).

The pharmaceutical compositions may be for human or animal usage in human and veterinary medicine and will typically comprise any one or more of a pharmaceutically acceptable diluent, carrier, or excipient. Acceptable carriers or diluents for therapeutic use are well known in the pharmaceutical art, and are described, for example, in
5 Remington's Pharmaceutical Sciences, Mack Publishing Co. (A. R. Gennaro edit. 1985). The choice of pharmaceutical carrier, excipient or diluent can be selected with regard to the intended route of administration and standard pharmaceutical practice. The pharmaceutical compositions may comprise as - or in addition to - the carrier, excipient or diluent any suitable binder(s), lubricant(s), suspending agent(s), coating
10 agent(s), solubilising agent(s).

Preservatives, stabilizers, dyes and even flavouring agents may be provided in the pharmaceutical composition. Examples of preservatives include sodium benzoate, sorbic acid and esters of p-hydroxybenzoic acid. Antioxidants and suspending agents
15 may be also used.

There may be different composition/formulation requirements dependent on the different delivery systems. By way of example, the pharmaceutical composition of the present invention may be formulated to be delivered using a mini-pump or by a mucosal
20 route, for example, as a nasal spray or aerosol for inhalation or ingestible solution, or parenterally in which the composition is formulated by an injectable form, for delivery, by, for example, an intravenous, intramuscular or subcutaneous route. Alternatively, the formulation may be designed to be delivered by both routes.

25 Where the pharmaceutical composition is to be delivered mucosally through the gastrointestinal mucosa, it should be able to remain stable during transit through the gastrointestinal tract; for example, it should be resistant to proteolytic degradation, stable at acid pH and resistant to the detergent effects of bile.

30 Where appropriate, the pharmaceutical compositions can be administered by inhalation, in the form of a suppository or pessary, topically in the form of a lotion, solution, cream, ointment or dusting powder, by use of a skin patch, orally in the form of tablets

containing excipients such as starch or lactose or chalk, or in capsules or ovules either alone or in admixture with excipients, or in the form of elixirs, solutions or suspensions containing flavouring or colouring agents, or they can be injected parenterally, for example intravenously, intramuscularly or subcutaneously. For parenteral
5 administration, the compositions may be best used in the form of a sterile aqueous solution which may contain other substances, for example enough salts or monosaccharides to make the solution isotonic with blood. For buccal or sublingual administration the compositions may be administered in the form of tablets or lozenges which can be formulated in a conventional manner.

ADMINISTRATION

Typically, a physician will determine the actual dosage which will be most suitable for an individual subject and it will vary with the age, weight and response of the particular
15 patient and severity of the condition. The dosages below are exemplary of the average case. There can, of course, be individual instances where higher or lower dosage ranges are merited.

The compositions (or component parts thereof) of the present invention may be
20 administered orally. In addition or in the alternative the compositions (or component parts thereof) of the present invention may be administered by direct injection. In addition or in the alternative the compositions (or component parts thereof) of the present invention may be administered topically. In addition or in the alternative the compositions (or component parts thereof) of the present invention may be
25 administered by inhalation. In addition or in the alternative the compositions (or component parts thereof) of the present invention may also be administered by one or more of: parenteral, mucosal, intramuscular, intravenous, subcutaneous, intraocular or transdermal administration means, and are formulated for such administration.

30 By way of further example, the pharmaceutical composition of the present invention may be administered in accordance with a regimen of 1 to 10 times per day, such as once or twice per day. The specific dose level and frequency of dosage for any

particular patient may be varied and will depend upon a variety of factors including the activity of the specific compound employed, the metabolic stability and length of action of that compound, the age, body weight, general health, sex, diet, mode and time of administration, rate of excretion, drug combination, the severity of the particular condition, and the host undergoing therapy.

The term "administered" also includes but is not limited to delivery by a mucosal route, for example, as a nasal spray or aerosol for inhalation or as an ingestible solution; a parenteral route where delivery is by an injectable form, such as, for example, an intravenous, intramuscular or subcutaneous route.

Hence, the pharmaceutical composition of the present invention may be administered by one or more of the following routes: oral administration, injection (such as direct injection), topical, inhalation, parenteral administration, mucosal administration, intramuscular administration, intravenous administration, subcutaneous administration, intraocular administration or transdermal administration.

DISEASES

Pharmaceutical compositions comprising an effective amount of a ScFv Ab and/or an NOI encoding same may be used in the treatment of disorders such as those listed in WO-A-98/09985. For ease of reference, part of that list is now provided: macrophage inhibitory and/or T cell inhibitory activity and thus, anti-inflammatory activity; anti-immune activity, i.e. inhibitory effects against a cellular and/or humoral immune response, including a response not associated with inflammation; diseases associated with viruses and/or other intracellular pathogens; inhibit the ability of macrophages and T cells to adhere to extracellular matrix components and fibronectin, as well as up-regulated fas receptor expression in T cells; inhibit unwanted immune reaction and inflammation including arthritis, including rheumatoid arthritis, inflammation associated with hypersensitivity, allergic reactions, asthma, systemic lupus erythematosus, collagen diseases and other autoimmune diseases, inflammation associated with atherosclerosis, arteriosclerosis, atherosclerotic heart disease,

reperfusion injury, cardiac arrest, myocardial infarction, vascular inflammatory disorders, respiratory distress syndrome or other cardiopulmonary diseases, inflammation associated with peptic ulcer, ulcerative colitis and other diseases of the gastrointestinal tract, hepatic fibrosis, liver cirrhosis or other hepatic diseases, thyroiditis or other glandular diseases, glomerulonephritis or other renal and urologic diseases, otitis or other oto-rhino-laryngological diseases, dermatitis or other dermal diseases, periodontal diseases or other dental diseases, orchitis or epididymo-orchitis, infertility, orchidial trauma or other immune-related testicular diseases, placental dysfunction, placental insufficiency, habitual abortion, eclampsia, pre-eclampsia and other immune and/or inflammatory-related gynaecological diseases, posterior uveitis, intermediate uveitis, anterior uveitis, conjunctivitis, chorioretinitis, uveoretinitis, optic neuritis, intraocular inflammation, e.g. retinitis or cystoid macular oedema, sympathetic ophthalmia, scleritis, retinitis pigmentosa, immune and inflammatory components of degenerative fundus disease, inflammatory components of ocular trauma, ocular inflammation caused by infection, proliferative vitreo-retinopathies, acute ischaemic optic neuropathy, excessive scarring, e.g. following glaucoma filtration operation, immune and/or inflammation reaction against ocular implants and other immune and inflammatory-related ophthalmic diseases, inflammation associated with autoimmune diseases or conditions or disorders where, both in the central nervous system (CNS) or in any other organ, immune and/or inflammation suppression would be beneficial, Parkinson's disease, complication and/or side effects from treatment of Parkinson's disease, AIDS-related dementia complex HIV-related encephalopathy, Devic's disease, Sydenham chorea, Alzheimer's disease and other degenerative diseases, conditions or disorders of the CNS, inflammatory components of strokes, post-polio syndrome, immune and inflammatory components of psychiatric disorders, myelitis, encephalitis, subacute sclerosing pan-encephalitis, encephalomyelitis, acute neuropathy, subacute neuropathy, chronic neuropathy, Guillain-Barre syndrome, Sydenham chorea, myasthenia gravis, pseudo-tumour cerebri, Down's Syndrome, Huntington's disease, amyotrophic lateral sclerosis, inflammatory components of CNS compression or CNS trauma or infections of the CNS, inflammatory components of muscular atrophies and dystrophies, and immune and inflammatory related diseases, conditions or disorders of the central and peripheral nervous systems, post-traumatic inflammation, septic shock,

infectious diseases, inflammatory complications or side effects of surgery, bone marrow transplantation or other transplantation complications and/or side effects, inflammatory and/or immune complications and side effects of gene therapy, e.g. due to infection with a viral carrier, or inflammation associated with AIDS, to suppress or inhibit a humoral and/or cellular immune response, to treat or ameliorate monocyte or leukocyte proliferative diseases, e.g. leukaemia, by reducing the amount of monocytes or lymphocytes, for the prevention and/or treatment of graft rejection in cases of transplantation of natural or artificial cells, tissue and organs such as cornea, bone marrow, organs, lenses, pacemakers, natural or artificial skin tissue. Specific cancer related disorders include but not limited to: solid tumours; blood born tumours such as leukemias; tumor metastasis; benign tumours, for example hemangiomas, acoustic neuromas, neurofibromas, trachomas, and pyogenic granulomas; rheumatoid arthritis; psoriasis; ocular angiogenic diseases, for example, diabetic retinopathy, retinopathy of prematurity, macular degeneration, corneal graft rejection, neovascular glaucoma, retrolental fibroplasia, rubeosis; Osler-Webber Syndrome; myocardial angiogenesis; plaque neovascularization; telangiectasia; hemophiliac joints; angiofibroma; wound granulation; coronary collaterals; cerebral collaterals; arteriovenous malformations; ischemic limb angiogenesis; neovascular glaucoma; retrolental fibroplasia; diabetic neovascularization; helicobacter related diseases, fractures, vasculogenesis, hematopoiesis, ovulation, menstruation and placentation.

Figures

The invention will now be further described only by way of example in which reference is made to the following Figures:

Figure 1 which shows a DNA sequence (SEQ ID No 5) encoding a 5T4 ScFv designated 5T4ScFv.1. The sequence of the mature secreted protein (SEQ ID No 1) is provided.

Figure 2 which shows a DNA sequence encoding a B7-1.5T4.1 fusion protein (SEQ ID No 7). A deduced amino acid sequence for the B7-1.5T4.1 fusion protein (SEQ ID No 3) is also provided.

5 Figure 3a which shows a diagrammatic representation of a B7-1.5T4.1 construct.

Figure 3b which shows a diagrammatic representation of a B7-1.5T4.2 construct.

10 Figure 4 which shows a DNA sequence encoding a B7-2.5T4.1 fusion protein (SEQ ID No 9). A deduced amino acid sequence for the B7-2.5T4.1 fusion protein (SEQ ID No 10) is also provided.

Figure 5 which shows a B7 link ScFv sequence (SEQ ID No 11).

15 Figure 6 which shows a DNA sequence encoding a Ig-5T4 fusion protein (SEQ ID No 8). A deduced amino acid sequence for the Ig-5T4 fusion protein (SEQ ID No 4) is also provided.

Figure 7 which shows an ScFv-IgE sequence (SEQ ID No 12).

20

Figure 8 which shows a B7-EGF sequence (SEQ ID No 13).

Figure 9 which shows the effect of the ScFv Ab on CT26-neo tumour cell growth in Balb/c mice over a period of 35 days.

25

Figure 10 which shows the effect of the ScFv Ab on CT26-h5T4 tumour cell growth in Balb/c mice over a period of 35 days.

30 Figure 11 which shows the effect of the ScFv Ab on B16-neo tumour cell growth in Balb/c mice over a period of 35 days.

Figure 12 which shows the effect of the ScFv Ab on B16-h5T4 tumour cell growth in Balb/c mice over a period of 35 days.

Figure 13 shows ScFv constructs.

5

Figure 14 shows a B7-scFv binding to the 5T4 target antigen.

Figure 15 shows B7-scFv binding to CTLA4.

10

Figure 16 shows FACS analysis of A9-5T4 (A) and A9-neo (5T4 negative) (B) cells incubated with scFv protein alone or scFv-HG1 fusion protein followed by goat anti-human IgG - FITC labelled antibody.

Figure 17 shows 5T4 scFv - Hy1 ADCC Activity.

15

In slightly more detail:

Figure 14 shows supernatants from mock transfected 293T cells or those transfected with the tagged B7-scFv construct were incubated with A9 5T4 and A9 neo cells.

20

Detection used FITC conjugated α His or α Myc antibodies.

Figure 15 shows A9 5T4 and A9 neo cells which were incubated with the scFv alone, a B7-scFv construct lacking the Myc-His tag or the tagged B7-scFv construct. The B7.1 ligand, CTLA4-Ig was added and detection used FITC conjugated α mouse IgG.

25

EXAMPLES

Example 1 – Construction of 5T4 ScFv Ab and retroviral - vector delivery to tumour.

30

The cDNA encoding the murine 5T4 monoclonal antibody is cloned and sequenced by standard techniques (Antibody engineering: a practical approach ed McCafferty *et al.* 1996 OUP). The sequence of the variable region of the antibody can be used to construct a ScFv antibody. The coding sequence of a 5T4 ScFv, called 5T4ScFv.1 (SEQ

ID No 1), is shown in Figure 1. In this molecule, the DNA sequence encodes the VH from the mouse 5T4 monoclonal antibody followed by a 15 amino acid flexible linker and the VL region of the mouse 5T4 antibody. The flexible linker encodes 3 copies of the amino-acid sequence gly-gly-gly-gly-ser and the DNA sequence similarity between the repeats has been minimised to avoid the risk of recombination between the repeats when plasmids containing them are grown in *E. coli*.

DNA CASSETTES

10 *Cassette 1 – Translation initiation signal and signal peptide*

In order to achieve correct translation initiation and secretion from mammalian cells, the following sequence is used:

15 aagcttCCACCATGGGATGGAGCTGTATCATCCTCTTCTTGGTAGCAACAGCTA
CAGGTGTCCACTCC

This contains a convenient *Hind*III restriction site for cloning into expression vectors (lower case), the consensus translation initiation signal for mammalian cells (ANNATGPu) and the coding sequence for a signal peptide sequence from an immunoglobulin gene.

Cassette 2 – scFv

25 The sequence of the secreted portion of the 5T4ScFv.1 is shown in Figure 1. This molecule can be represented as Vh – (gly₄-ser)₃ linker - Vl.

The 5T4 ScFv2 Ab consists of the 5T4 variable region sequences connected in the order Vl – flexible linker Vh. In this case the linker encodes the 20 amino-acid peptide (gly₄-ser)₄. A longer linker improves assembly of the ScFv when the V-region segments are in this order. (Pluckthun *et al* in Antibody Engineering: a practical approach, ed McCafferty *et al*. 1996 OUP).

EXPRESSION OF A 5T4 SPECIFIC SCFV

For expression of a 5T4-specific ScFv in human cells, the coding sequence is inserted into the vector pCIneo (Promega) under the control of a strong promoter and polyadenylation signal. The translation initiation signal and immunoglobulin leader (signal peptide) sequence from Cassette 1 at the 5'end of the coding region ensure efficient secretion of the ScFv from mammalian cells.

Example 2 – Transfection of macrophages/monocytes with an expression vector encoding an ScFv Ab.

Peripheral blood mononuclear cells are isolated from human peripheral blood at laboratory scale by standard techniques procedures (Sandlie and Michaelsen 1996 In Antibody engineering: a practical approach. Ed McCafferty *et al.* Chapter 9) and at large scale by elutriation (eg Ceptra from CellPro). Adherent cells (essentially monocytes) are enriched by adherence to plastic overnight and cells can be allowed to differentiate along the macrophage differentiation pathway by culturing adherent cells for 1-3 weeks.

Monocytes and macrophages are transfected with an expression vector capable of expressing a ScFv Ab in human cells. For constitutive high level expression, the ScFv Ab is expressed in a vector which utilises the hCMV-MIE promoter-enhancer, pCI (Promega). For hypoxia-induced expression, the hCMV promoter is replaced by a promoter containing at least one HRE. A suitable promoter is a truncated HSV TK promoter with 3 copies of the mouse PGK HRE (Firth *et al.* 1994 Proc. Natl. Acad. Sci. 91: 6496-6500).

A variety of transfection methods can be used to introduce vectors into monocytes and macrophages, including particle-mediated DNA delivery (biolistics), electroporation, cationic agent-mediated transfection (eg using Superfect, Qiagen). Each of these methods is carried out according to the manufacturer's instructions, taking into account the parameters to be varied to achieve optimal results as specified by the individual

manufacturer. Alternatively, viral vectors may be used such as defective Adenovirus vectors (Microbix Inc or Quantum Biotechnologies Inc).

Example 3 - Construction of B7 – ScFv Fusion proteins

5

The extracellular domain of B7-1 is defined by amino-acid residues 1 – 215 of the native human B7-1 protein. This sequence, together with its signal peptide-encoding sequence, is used to construct secreted fusion proteins which also contain the ScFv derived from the 5T4 monoclonal antibody. The sequence of the 5T4 ScFv is given in
10 Figure 1.

A DNA coding sequence is constructed using standard molecular biology techniques which encodes a fusion protein in which the N-terminus of the 5T4 ScFv is fused after amino acid 215 of human B7-1. The sequence of this coding sequence, B7-1.5T4.1
15 (SEQ ID No 7) is shown in Figure 2. The fusion protein contains a flexible (gly-gly-gly-gly-ser) spacer between the B7-1 and 5T4 ScFv sequences. The introduction of a convenient BamH1 restriction site at the end of the linker insertion (beginning at nucleotide 733) also allows for further linkers to be screened for optimal expression of bi-functional fusion protein. Figure 3 indicates the fusion protein in diagrammatic
20 form. It is similarly possible to construct B7-1.5T4.2 (Figure 3b) in which the ScFv is N-terminal and the B7 extracellular domain is C-terminal. In this case only the coding sequence of the mature B7-1 (without signal peptide) is required. A signal peptide such as an immunoglobulin leader sequence is added to the N-terminus of the ScFv in this instance.

25

For fusion proteins which use the co-stimulatory extracellular domain of B7-2 (Gerstmayer et al 1997 J Immunol 158(10): 4584-90), the signal peptide and extracellular domain of B7-2 is used in place of B7-1 sequences. Figure 4 shows the coding sequence of the SCM B7-2.5T4.1co-stimulatory domain. It encodes the first
30 225 amino acids of human B7-2, preceded by its signal peptide, and a flexible linker (gly4-ser). The BamHI site at the end of this sequence can be used to insert the domain upstream of the 5T4ScFv.1 (see Figure 3). The sequence includes the B7-2 signal

peptide which can serve to allow secretion of this fusion protein in which the B7-2 domain is at the N-terminus of the fusion protein.

Each engineered cDNA is inserted into the mammalian expression vector pCI to allow
 5 expression in mammalian tissue culture cells. For this purpose, a linker sequence is added to the 5'-end of the coding sequence which introduces a convenient restriction site for insertion into the polylinker of pCI and adds the translation initiation signal CCACC immediately adjacent to the first ATG codon. Constructs in pCI are transfected into a suitable mammalian host cell line such as COS-1 to confirm secretion
 10 of the SCM. The transcription cassette from pCI or an appropriate segment of the transcription cassette is subsequently sub-cloned into the expression vector to be used as the gene delivery system for therapeutic use.

**Example 4 - Transfection of macrophages/monocytes with an expression vector
 15 encoding a ScFv Ab comprising a secreted co-stimulatory molecule (SCM).**

Peripheral blood mononuclear cells are isolated from human peripheral blood at laboratory scale by standard techniques procedures (Sandlie and Michaelsen 1996 In Antibody engineering: a practical approach. Ed McCafferty *et al.* Chapter 9) and at
 20 large scale by elutriation (eg Cephate from CellPro). Adherent cells (essentially monocytes) are enriched by adherence to plastic overnight and cells can be allowed to differentiate along the macrophage differentiation pathway by culturing adherent cells for 1 – 3 weeks.

25 Monocytes and macrophages are transfected with an expression vector capable of expressing an ScFv Ab comprising an SCM in human cells. For constitutive high level expression, the SCM is expressed in a vector which utilises the hCMV-MIE promoter-enhancer, pCI (Promega). For hypoxia-induced expression, the hCMV promoter is replaced by a promoter containing at least one HRE. A suitable promoter is a truncated
 30 HSV TK promoter with 3 copies of the mouse PGK HRE (Firth *et al.* 1994 Proc. Natl. Acad. Sci. 91: 6496-6500).

A variety of transfection methods can be used to introduce vectors into monocytes and macrophages, including particle-mediated DNA delivery (biolistics), electroporation, cationic agent-mediated transfection (eg using Superfect, Qiagen). Each of these methods is carried out according to the manufacturer's instructions, taking into account the parameters to be varied to achieve optimal results as specified by the individual manufacturer. Alternatively, viral vectors may be used such as defective Adenovirus vectors (Microbix Inc or Quantum Biotechnologies Inc).

Example 5 - Analysis of SCM binding to CTLA-4 and 5T4-antigen expressing cells

10

The B7-1 or B7-2 domains of an ScFv Ab-SCM fusion protein are expected to bind specifically to CD28 and CTLA-4 present on human T-cells. Binding to T-cells or Chinese hamster ovary cells transfected with human CTLA-4 or CD28 is determined using FACS analysis as follows. 5×10^5 CTLA-4 expressing target cells or equivalent cells lacking CTLA-4 (untransfected CHO cells) are incubated with 0.1 ml culture supernatant from COS-1 cells transiently transfected with SCM genes for 1 h at 4°C. The cells are washed and incubate with 1 mg monoclonal antibody specific for the B7 domain (eg Mab 9E10) followed by FITC-labelled goat anti-mouse IgG (Pharmingen) and analysis by FACS.

20

Binding of ScFv to 5T4-antigen is similarly assessed using target cells expressing 5T4-antigen (5T4-transfected A9 cells) or control cells (A9).

Example 6 - Analysis of co-stimulatory activity

25

An established mouse cell line of Balb/c origin such as HC11 cells is transfected with the cDNA encoding human 5T4-antigen (Myers *et al.* 1994 J. Biol. Chem. 269; 9319-9324) inserted in the expression vector pCIneo.

30

Splenic T-cells from Balb/c mice are isolated by standard procedures (Johnstone and Thorpe 1996 In Immunochimistry in Practice. Blackwell. Chapter 4). T-cells are pre-stimulated by incubation for 1 – 2 days in medium containing 10ng/ml PMA (Sigma)

and 100 U/ml human IL-2 (Boehringer Mannheim). HC11-5T4 cells are incubated at 10^4 cells /well of a 96-well tissue culture tray for 2 h with up to 0.1ml supernatant from COS cells transfected with SCM gene. Up to 10^5 pre-stimulated T-cells are added to each well, the cells are pulsed with 0.25 mCi / well ^3H -thymidine and incorporation of ^3H -thymidine is measured using a liquid scintillation counter after 24h.

Example 7 - Analysis of co-stimulation in animal models.

HC11 cells transfected with the human 5T4-antigen gene are grown as tumours in Balb/c mice. SCM genes B7-1.5T4.1 or B7-2.5T4.1 or a combination of both genes are introduced into the tumour cells prior to implantation and the growth of the tumours and the growth of control tumours which do not express SCM genes *in vivo* are monitored.

Example 8 - Construction of a B7-1/ScFv, specific for human 5T4, fusion protein

Standard molecular biology techniques are used to construct a fusion protein consisting of the leader sequence and extracellular domain of B7-1, fused via a flexible linker to the V_H and V_L of the murine Mab 5T4 specific to human 5T4.

The flexible linker, used to join the extracellular domain of B7.1 and the ScFv, was constructed by annealing two homologous oligonucleotides with engineered 5' Sma I and 3' Spe I sites - using oligonucleotides

upper

5' GGG GGT GGT GGG AGC GGT GGT GGC GGC AGT GGC GGC GGC GGA A
3'

and lower

5' CTA GTT CCG CCG CCG CCA CTG CCG CCA CCA CCG CTC CCA CCA CCC
CC 3'

The linker is cloned into pBluescript (Stratagene) via Sma I and Spe I to produce pLINK. The signal peptide (sp) and extracellular domain of murine B7.1 were amplified by PCR from pLK444-mB7.1 (supplied by R. Germain NIH, USA) via primers that introduce 5' EcoRI and 3' Sma I sites - primers forward

5

5' C TCG AAT TCC ACC ATG GCT TGC AAT TGT CAG TTG ATG C 3'

reverse

10

5' CTC CCC GGG CTT GCT ATC AGG AGG GTC TTC 3'

The B7.1 PCR product was cloned into pLINK via Eco RI and Sma I to form pBS/B7Link.

15

The V_H and V_L of the 5T4 specific ScFv was amplified via primers -

forward primer

5' CTC ACT AGT GAG GTC CAG CTT CAG CAG TC 3'

20

reverse primer

5' CTC GCG GCC GCT TAC CGT TTG ATT TCC AGC TTG GTG CCT CCA CC 3'

25

that introduce 5' Spe I and 3' Not I sites from pHEN1-5T4 ScFv. PBS/B7Link was digested with Spe I and Not I and ligated with the ScFv to form OBM 233 consisting of the sequence shown as SEQ ID No. 11: B7 Link ScFv sequence (Figure 5).

30

This fusion can be used to construct a recombinant vector e.g. retrovirus, Lentivirus, adenovirus, poxvirus, vaccinia virus, baculovirus. Such vectors can be used to inject patient tumours directly. To deliver the fusion protein to tumour cells the recombinant vector is used to transduce macrophages/monocytes/CD34+ cells *ex vivo* before

injection back into patients. These cells will traffic to tumours. The ScFv will bind to a specific tumour antigen expressed on the surface of tumour cells e.g. 5T4 (Myers et al 1994 JBC). B7 is found on the surface of professional antigen presenting cells e.g. macrophages, dendritic cells and B cells. It interacts with its ligands CD28 and CTLA-4 located on CD4 and CD8 cells. The simultaneous interaction of B7-CD28/CTLA-4 and MHC-peptide/T cell receptor leads to a pronounced increase in IL-2 which promotes CD8 (cytotoxic T cell) expansion (Linsley PS, Brady W, Grosmaire L, Aruffo A, Damle NK, Ledbetter JA J Exp Med 1991 Mar 1;173(3):721-730 Binding of the B-cell activation antigen B7 to CD28 costimulates T cell proliferation and IL-2 mRNA accumulation.) Tumour cells that have been B7 transfected with B7 have been shown retardation in animal models (Townsend SE, Allison JP Science 1993 15;259(5093):368-370).

Example 9 - Transient expression and purification of B7-1/ScFv and Leader ScFv (LScFv)

For transient expression of B7-1/ScFv the human CMV expression plasmid pCIneo (Promega) was used. B7/ScFv was excised from OBM 233 by digestion with EcoRI/Not I and cloned into pCIneo that was previously digested with EcoRI/Not I. Transient expression of recombinant protein is made by transfection of 293T cells with the relevant plasmid using calcium phosphate (Profectin, Promega). Conditions used were similar to those recommended by the manufacturer. To reduce bovine serum contamination serum free optimum media (Gibco BRL). After 36-48 hours transfection supernatants were harvested and spun through a Centrprep (Amicon, Glos. UK) 10 filter (all proteins larger than 10 kDa are purified/concentrated) and a Centricon (Amicon) 10 filter. Supernatants are concentrated approximately 30 fold.

For B7-1 to be biologically functional it must be able to display binding with one of its natural ligands either CTLA-4 or CD28 found on the surface of specific populations of T cells (e.g CD4+). The biological activity B7-1/ScFv fusion protein was analysed for simultaneous interaction with its natural ligand CTLA-4 (in the form of CTLA4-Ig supplied by Ancell, MN, USA) and A9 cells expressing human 5T4. Briefly:

approximately 5×10^5 A9-h5T4 cells were incubated with 100ul of either B7.1/ScFv or LScFv supernatant in a U bottom 96 well plate at 4°C for 1 hour. After washing cells were incubated with CTLA4-Ig (Ancell) for 1 hour. After washing, bound CTLA4-Ig was detected using an FITC conjugated anti-mouse Ig (Dako).

Results show obvious binding of CTLA-Ig with the B7-1 extracellular domain, bound via the ScFv, to the surface of human 5T4 positive A9 cells. The lack of binding activity with 5T4 negative A9 cells further illustrates that the interaction of B7 with CTLA4-Ig and ScFv with 5T4 are specific.

Example 10 - ScFv-IgG fusion example

Construction of ScFv-IgG

The sequence encoding a translation initiation sequence and the human immunoglobulin kappa light chain signal peptide is synthesised as two complementary single stranded oligonucleotides which when annealed also contain an internal *Xho I* site at the 5' end and in addition leave a *Xba I* compatible 5' overhang and a *Pst I* compatible 3' overhang

ctagactcgagCCACC ATG GGA TGG AGC TGT ATC ATC CTC TTC TTG GTA
GCA ACA GCT ACA GGT GTC CAC TCC GAG GTC CAG ctgca

and

g CTG GAC CTC GGA GTG GAC ACC TGT AGC TGT TGC TAC CAA GAA GAG
GAT GAT ACA GCT CCA TCC CAT GGTGGctcgagt

This is then cloned into pBluescript II (Stratagene) restricted with *Xba I* and *Pst I* to create pBSII/Leader.

The 5T4 ScFv is amplified by PCR from pHEN1 using oligonucleotides which incorporate a *Pst I* site at the 5' end of the product and a *Hind III* at the 3' end

GTC CAG CTG CAG CAG TCT GG

5

and

CG TTT GAT TTC AAG CTT GGT GC

10 This is then restricted with those enzymes and inserted into pBSII/Leader restricted with the same enzymes, creating pBSII/Leader/ScFv.

The HIgG 1 constant region is amplified by PCR from the cloned gene using oligonucleotides which incorporate a *Hind III* site at the 5' end and a *Xho I* site at the 3'

15

end

gcgc AAG CTT gaa atc aaa cgg GCC TCC ACC AAG GGC CCA

and

20

gcgc ctcgag TCA TTT ACC CGG AGA CAG GG

This is then restricted with those enzymes and inserted into pBSII/Leader/ScFv restricted with the same enzymes, creating pBSII/Leader/ScFv/HG1. The sequence for this construct is shown in the Figure 4 (SEQ ID No 10).

25

This fusion can be used to construct a recombinant vector e.g. retrovirus, Lentivirus, adenovirus, poxvirus, vaccinia virus, baculovirus. Such vectors can be used to inject patient tumours directly. To deliver the fusion protein to tumour cells the recombinant vector is used to transduce macrophages/monocytes/CD34+ cells *ex vivo* before
30 injection back into patients. These cells will traffic to tumours. The ScFv will bind to a specific tumour antigen expressed on the surface of tumour cells e.g. 5T4 (Myers et al 1994 JBC). Bound IgG will promote specific tumour destruction via a collection of

mechanisms collectively known as antibody dependent cellular cytotoxicity (Munn et al Can res 1991 *ibid*, Primus et al 1993 Cancer Res *ibid*).

Example 11 - Construction of ScFv-IgE1 (human IgE1 heavy constant region)

5

A similar fusion construct of 5T4 ScFv – human IgE constant heavy chain is made consisting of the sequence shown as Figure 7 (SEQ ID No. 12).

This fusion construct is made by amplifying the human IgE1 constant heavy region by
10 PCR cDNA derived from human B-cells RNA by RT and subsequently using oligonucleotides which incorporate a *Hind III* site at the 5' end and a *Xho I* site at the 3' end

gcgc AAG CTT gaa atc aaa cgg GCC TCC ACA CAG AGC CCA

15

and

gcgc ctcgag TCA TTT ACC GGG ATT TAC AGA

20 This is then restricted with those enzymes and inserted into pBSII/Leader/ScFv restricted with the same enzymes, creating pBSII/Leader/ScFv/HE1.

As described above the ScFv-IgE construct can be incorporated into a recombinant viral vector for use in gene therapy of cancer e.g. inject patient tissue directly or to transduce
25 patient derived macrophages/monocytes/CD34+ cells *ex vivo*. The fusion protein will be secreted and will bind to tumour cells bearing the antigen that the ScFv is specific for. Binding of IgE to tumour cells should promote a strong histamine response via activation of mast cells. This will lead to a strong inflammatory response and destruction tumour cells as is reported for IgE cytotoxic destruction of parasites e.g.
30 helminth larvae (Capron M 1988 Eosinophils in diseases: receptors and mediators. In progress in allergy and clinical immunology (Proc. 13th Int. Congress of Allergy and Clinical Immunology) Hogrefe & Huber Toronto p6). Such inflammation and tumour

destruction should initiate the recruitment of other immune effector cells. Past reports indicate that treatment with an MMTV antigen specific IgE Mab leads to protection from a tumour expressing MMTV antigen (Nagy E, Istavvan B, Sehon AH 1991 Cancer Immunol. Immunotherapy vol 34:63-69).

5

Example 12 - Construction of B7/EGF

B7 – EGF Synthetic Gene.

10 A fusion construct of B7 – EGF is made by inserting a PCR product amplified from the region of the gene encoding the mature EGF peptide (see accession number X04571) into pBS/B7 Link. This construct has the sequence shown in Figure 8 (SEQ ID No. 13).

15 Using cDNA derived by RT of RNA isolated from a cell line such the 293 human kidney line (ATCC: CRL1573), the DNA is amplified by PCR using oligonucleotides containing a *Spe I* restriction enzyme site at the N-terminus and a stop codon and a *Not I* site at the C-terminus

20 GG ACT AGT AAT AGT GAC TCT GAA TGT CCC

And

ATT AGC GGC CGC TTA GCG CAG TTC CCA CCA CTT C

25

The resulting product is digested with those enzymes and ligated to pBS/B7 Link which has been restricted with the same enzymes creating pBS/B7 Link EGF. The B7 Link EGF cassette is then excised with *Eco RI* and *Not I* and inserted into a derivative of pHIT111 (Soneoka *et al*, 1995, Nucl Acid Res 23; 628) which no longer carries the

30 *LacZ* gene

An alternative to using ScFv is to use growth factors that have a high affinity to their corresponding receptor e.g. epidermal growth factor which binds to several receptors including erb-2 which is highly associated with tumourgenesis.

- 5 As described above the fusion construct can be incorporated into a recombinant viral vector for use in gene therapy e.g. inject patient tissue directly or to transduce patient derived macrophages/monocytes/CD34+ cells *ex vivo*. The fusion protein will be secreted and will bind to tumour cells bearing the erb-2 antigen.
- 10 Epidermal growth factor (EGF) will bind to its ligand erb-2 (an EGF receptor) thus obviating the requirement of a ScFv. Erb-2 is highly associated with tumour cells (Hynes NE Semin Cancer Biol 1993 Feb;4(1):19-26, Amplification and over expression of the erbB-2 gene in human tumors: its involvement in tumor development, significance as a prognostic factor, and potential as a target for cancer therapy). B7 is
15 found on the surface of professional antigen presenting cells e.g. macrophages, dendritic cells and B cells. It interacts with its ligands CD28 and CTLA-4 located on CD4 and CD8 cells. The simultaneous interaction of B7-CD28/CTLA-4 and MHC-peptide/T cell receptor leads to massive increase in IL-2 which promotes CD8 (cytotoxic T cell) expansion (Linsley PS, Brady W, Grosmaire L, Aruffo A, Damle NK, 20 Ledbetter JA J Exp Med 1991 Mar 1;173(3):721-730 Binding of the B cell activation antigen B7 to CD28 costimulates T cell proliferation and interleukin 2 mRNA accumulation.) Tumour cells that have been B7 transfected with B7 have shown retardation in animal models (Townsend SE, Allison JP Science 1993 15;259(5093):368-370 Tumor rejection after direct costimulation of CD8+ T cells by 25 B7-transfected melanoma cells). It has been reported that B7 will enhance the CTL response to tumour antigens specific to tumour cells thus leading to the destruction of all such cells.

Example 13 - Production of cell lines expressing fusion constructs

30

The ScFv-IgG gene was excised from pBSII/L/ScFv/hIgG1 by Xho I digestion, and cloned into pLXSN via the Xho I site, to make pLXSN/ScFv-IgG, such that after

chromosomal integration it is under transcriptional control of the LTR. Virus was made in the human kidney cell line 293T by co-transfecting plasmids containing the MLV gap-pol genes (pCIEGPPD) and the VSV G envelope (pRV67) using the triple plasmid HIT system (Landau & Littman 1992 J Virol 66 5110, Soneoka Y et al 1995 NAR 23:628-633). Virus is harvested after 48 hours and used to transduce BHK-21 cells (ATCC# CCL-10). Approximately 24 hours post-transduction, transduced cells are selected by the addition of 1mg/ml G418 (Gibco BRL) to culture medium. The supernatant from positive colonies was harvested and concentrated by centrifugation through a Centriprep (Amicon, Glos. UK) 10 filter (all proteins larger than 10 kDa are purified/concentrated) and a Centricon (Amicon) 10 filter. Supernatants were concentrated approximately 30 fold.

Other fusion proteins are cloned into pLXSN via the Xho I site and expressed and concentrated using a similar protocol.

FACS analysis of fusion protein binding with cells expressing specific ligand

To determine if the ScFv-IgG fusion protein is specific for its antigen, human 5T4, FACS analysis of a human bladder carcinoma tumour line (EJ) or a stable murine cell line expressing h5T4, A9-h5T4 (Myers et al 1994 JBC) and a 5T4 negative line A9-neo was carried out. Approximately 5×10^5 A9 or EJ cells, in a round bottom 96 well plate (Falcon) were incubated with 100ul of a 1:5 dilution of concentrated supernatant (as described above) for 1 hour at 4°C. After washing, bound protein is detected using an anti human IgG/FITC conjugated antibody (Dako). Cells were analysed on a Becton Dickinson FACS machine. FACS results show that there is at least a 1 log shift in fluorescence activity in those 5T4 positive cells treated with the ScFv-IgG construct compared to the negative control construct consisting of the ScFv protein alone. A9 neo FACS shows that there is no non-specific binding of the ScFv component of the fusion protein.

FACS analysis of ScFv-IgE is carried out similar to above except that anti-human IgE-FITC (Dako) is used to detect binding of the fusion protein.

The B7/EGF fusion protein is analysed for binding using FACS and HC11-erb-2 positive cells (Hynes et al 1990). CTLA4-Ig (Ancell, USA) is used to analyse the bioactivity of the B7 component of the bound fusion protein. Anti-mouse IgG-FITC is used to show CTLA-4 binding.

5

Analysis of Fusion Proteins

Facs analysis of B7-scFv

10 Recombinant protein was generated by expression from a stably transfected BHK-21 cell line as described below (to allow identification and also purification) c-terminal to the scFv (Figure 13B) in the plasmid pCIneo (Promega) . To demonstrate that the scFv is able to bind to the 5T4 antigen, supernatants from these and from mock transfected 293T cells were added to mouse A9 cells expressing h5T4 (stable transfectants with a
15 h5T4 / neomycin resistance expression construct). Detection using FITC conjugated α His or α Myc antibodies confirmed binding of the scFv to the A9-5T4 cells but not 5T4 negative A9 neo cells indicating that the fusion construct is able to bind the target antigen (Figure 14).

20 Further FACS analysis was undertaken to show that the B7-scFv protein is able to bind simultaneously the B7.1 ligand, CTLA4 and cells expressing h 5T4. A9 5T4 and A9 neo cells were incubated with the scFv alone, a B7-scFv construct lacking the Myc-His tag or the tagged B7-scFv construct. The B7.1 ligand, CTLA4-Ig was added and detected using FITC conjugated α mouse IgG (Figure 15). The presence or absence of
25 the Myc-His tag made little difference to the simultaneous binding of the protein to 5T4 antigen and CTLA-4.

Analysis of 5T4 scFv HIgG 1 protein

30 Recombinant protein was generated by stable transfection of BHK-21 cells with constructs containing either 5T4 scFv alone or 5T4scFv-Hg1 (Figure 13A & C respectively) fusion under the control of the CMV immediate / early promoter. Figure 16 shows FACS analysis of mouse A9 5T4 cells. The cells were incubated with cell

supernatant from BHK-21 cells expressing either scFv alone or scFv-HG1, followed by goat anti-human IgG - FITC labelled antibody. As can be seen the scFv-HG1 is able to bind the 5T4 expressing cells and can be detected with the goat anti-human IgG - FITC labelled antibody. Figure 16b shows that this is due to the presence of 5T4 at the cell surface since no binding is observed with A9 cells that express the neomycin resistance marker, but no h5T4.

The same supernatants were used in an antibody dependent cell-mediated cytotoxicity (ADCC) assay which demonstrated that the scFv-Hy1 fusion protein is able to direct lysis of A9 5T4 cells. The A9 5T4 and neo cell lines were used in a chromium release assay. After labelling with ^{51}Cr , cells were incubated with either no protein the scFv alone or the scFv-Hy1 fusion construct. Freshly isolated peripheral blood lymphocytes were added and incubated for 4 hours. An aliquot of supernatant was taken for scintillation counting.

15

% lysis was calculated as :
$$\frac{\text{Test Release} - \text{Spontaneous Release}}{\text{Maximum Release} - \text{Spontaneous Release}} \times 100$$

Up to ~40% lysis was obtained with increasing effector:target ratio when compared to the scFv alone. The 5T4 negative cell line showed no increased lysis (Figure 17).

20

Example 14 – Analysis of efficacy in animal models

Human tumour-derived cell lines and tissues are cultured *in vivo* in genetically immunodeficient, “nude” mice according to well established techniques (see for example Strobel *et al.* 1997 Cancer Res. 57: 1228-1232; McLeod *et al.* 1997 Pancreas 14: 237-248). Syngeneic mouse models, in which a syngeneic tumour line is introduced into an immunocompetent mouse strain may also be used. These serve as suitable animal models for evaluating gene delivery systems of the invention. Vectors or engineered cells are administered systemically or directly into the tumour and tumour growth is monitored in treated and untreated animals. This system is used to define the

30

effective dose range of the treatments of the invention and the most appropriate route of administration.

ScFv Fusion Protein *In vivo* Anti Tumour Efficacy Data

5

The purpose of the study was to test the efficacy of a series of single chain antibody fusion proteins.

Murine mouse models, based on CT26, a chemically induced adenocarcinoma of
10 BALB/c origin (Brittain *et al.*, (1980) Cancer Res. 40:179-184), and on B16, a melanoma line derived from C57 B6 mice. Both the CT26 line and B16 are stably transformed to express human and murine 5T4. Mice are injected I.V. (to induce lung nodules, CT26) or subcutaneously (CT26 and B16) to make single mass subcutaneous tumours.

15

Experimental Design

CT26 cells expressing human 5T4 (CT26-h5T4) and CT26-neo

20

Cells were pre-incubated with:

PBS, LScFv-1, LScFv-2, B7-ScFv, ScFv-Ig

LScFv-1 and 2 were expressed in a BHK cell line. LScFv-1 was purified via its
25 Histidine tag on a Nickel column and ScFv-2 was purified using a filtration system. B7-ScFv was purified from a BHK line via a His tag and ScFv-Ig was purified via a filtration column. The concentration of each ScFv used in the experiment was defined as the amount of protein required to saturate binding of CT26-h5T4 cells in a FACS assay.

30

CT26-h5T4 and CT26-neo cells were pre-incubated with saturating amounts of each ScFv and incubated for 1 hour. After washing cells 5×10^5 cells were injected subcutaneously into the flanks of syngeneic BALB/c mice.

5 Tumour measurements were taken every two days and the volume calculated.

Results 14

Figure 9: CT26-neo

10

There is not a significant difference between the groups studied apart from the treatment with LScFv-1, for which there is an approximate 3-fold reduction in tumour size compared to the PBS control 36 days after tumour inoculation.

15 Figure 10: CT26-h5T4

Tumours treated with all of the 5T4 ScFv constructs had a significant effect on tumour growth. Four of the five mice treated with 5T4 ScFv-1 were tumour free on day 36. On day 36 ScFv-1 treated tumour cells were >60 fold smaller than tumours treated with
20 PBS.

When similar experiments were carried out using a mouse melanoma line (B16) engineered to express h5T4 a minimal anti-tumour effect was found with the ScFv constructs used (see Figures 11 and 12). The CT26 appear to be more sensitive to anti-
25 tumour immune responses induced by ScFv binding than the B16 cells. In addition, B16 cells do not express murine 5T4 whereas CT26 cells have mRNA to murine 5T4.

In summary there appears to be no benefit of fusing B7 or IgG to the 5T4 specific ScFv in the CT26 and B16 murine models. In fact, we have found in our examples that the
30 ScFv alone is more efficacious than the ScFv fusion constructs due to its higher binding affinity (as shown in BIACORE compared to B7-ScFv). Therefore these data indicate that the ScFv alone has a significant effect on tumour retardation and immune

enhancing molecules fused to the ScFv may not be required to show an effect on tumour retardation in the 5T4 model .

Summary

5

The present invention therefore provides antibodies capable of recognising a disease associated cell surface marker (DACSM). These antibodies may be used in the diagnosis and treatment of diseases associated with a DACSM.

10 All publications mentioned in the above specification are herein incorporated by reference. Various modifications and variations of the described methods and system of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the
15 invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be covered by the present invention.

20

CLAIMS

1. Use of an ScFv Ab (ScFv Ab) capable of recognising a disease associated cell surface molecule (DACSM) in the manufacture of a medicament for the prevention and/or treatment of a disease condition associated with a DACSM.
5
2. Use of an ScFv Ab according to claim 1 wherein the DACSM is a tumour associated antigen (TAA).
- 10 3. Use according to claim 1 or claim 2 wherein the ScFv Ab has the sequence presented as SEQ ID No 1 or SEQ ID No 2 or a variant, homologue, fragment or derivative thereof.
4. Use according to claim 1 or claim 2 wherein the ScFv Ab has the sequence presented as SEQ ID No 3 or a variant, homologue, fragment or derivative thereof.
15
5. Use according to claim 1 or claim 2 wherein the ScFv Ab has the sequence presented as SEQ ID No 4 or a variant, homologue, fragment or derivative thereof.
- 20 6. A nucleotide sequence encoding the ScFv Ab according to any one of claims 1-5.
7. A nucleotide sequence according to claim 6 wherein the nucleotide sequence has the sequence presented as SEQ ID No 5 or SEQ ID No 6 or a variant, homologue, fragment or derivative thereof.
25
8. A nucleotide sequence according to claim 6 wherein the nucleotide sequence has the sequence presented as SEQ ID No 7 or a variant, homologue, fragment or derivative thereof.
30
9. A nucleotide sequence according to claim 6 wherein the nucleotide sequence has the sequence presented as SEQ ID No 8 or a variant, homologue, fragment or

derivative thereof.

10. A nucleotide sequence capable of hybridising to the nucleotide sequence according to any one of claims 6-9 or a sequence that is complementary to the hybridisable nucleotide sequence.

11. A nucleotide sequence according to any one of claims 6-10 wherein the nucleotide sequence is operably linked to a promoter.

12. A construct comprising the nucleotide sequence according to any one of claims 6-11.

13. A vector comprising the nucleotide sequence of any one of claims 6-12.

14. A plasmid comprising the nucleotide sequence of any one of claims 6-13.

15. A host cell comprising the nucleotide sequence of any one of claims 6-14.

16. A process for preparing a ScFv Ab according to any one of claims 1-5 comprising expressing a nucleotide sequence according to any one of claims 6-11 or when present in the expression entity of any one of claims 12-15 and optionally isolating and/or purifying the ScFv Ab.

17. A ScFv Ab produced by the process according to claim 16.

18. An *in vitro* method for obtaining a ScFv Ab according to any of the preceding claims comprising:

(i) preparing a phage library wherein each phage comprises a nucleic acid construct encoding a protein comprising a potential binding domain;

- (ii) causing the expression of said potential proteins and the display of the potential binding domains on the outer surface of the phage;
 - (iii) contacting the phage library with a DACSM target under conditions such that the potential binding domains and the DACSM target interact;
 - (iv) separating the phage displaying a domain that binds the DACSM target from phage that do not bind;
 - (v) recovering at least one phage displaying on its outer surface a protein which binds the DACSM target;
 - (vi) amplifying the binding protein *in vitro* to create a second enriched library of binding structures;
 - (vii) repeating steps (iii) to (vi) at least twice;
 - (viii) expressing the nucleic acid encoding the binding protein under *in vitro* conditions; and
 - (ix) determining whether the binding protein interacts with the DACSM by detecting the presence or absence of a signal.
19. An *in vitro* method according to claim 18 wherein the *in vitro* method is to screen for a ScFv Ab useful in the treatment of a disease.
20. A process comprising the steps of:
- (a) performing the *in vitro* method according to claim 18 or claim 19;
 - (b) identifying one or more ScFv Abs capable of recognising a DACSM by means of a detectable signal; and

- (c) preparing a quantity of those one or more ScFv Abs.

21. A process comprising the steps of:

5 performing the method according to claim 18 or claim 19;

- (b) identifying one or more ScFv Abs capable of recognising a DACSM by means of a detectable signal; and

10 (c) preparing a pharmaceutical composition comprising those one or more identified ScFv Abs.

22. A process comprising the steps of:

15 (a) performing the method according to claim 18 or claim 19;

- (b) identifying one or more ScFv Abs capable of recognising a DACSM;

(c) modifying those one or more identified ScFv Abs capable of recognising a
20 DACSM; and

- (d) preparing a pharmaceutical composition comprising those one or more modified ScFv Abs.

25 23. A ScFv Ab as defined in any one of claims 1 to 5 or according to claim 17 or identified by the *in vitro* method of claim 18 or 19 wherein the ScFv Ab is capable of recognising a TAA.

24. A ScFv Ab according to claim 23 wherein the ScFv Ab is capable of recognising
30 a 5T4 antigen.

25. A method of affecting a disease *in vivo* with an ScFv Ab; wherein the ScFv Ab recognises a DACSM antigen in an *in vitro* method; and wherein the *in vitro* method is the method defined in claim 18 or claim 19.

5 26. Use of a ScFv Ab as defined in any one of claims 1 to 5 as defined in claim 17 or claim 23 or claim 24 to prepare a pharmaceutical composition.

27. A pharmaceutical composition as defined in claim 26 comprising a ScFv Ab and another therapeutically useful agent.

10

28. A pharmaceutical composition according to claim 27 wherein the other therapeutically useful agent is a pro-drug activating enzyme.

15 29. A pharmaceutical composition according to claim 28 wherein the other therapeutically useful agent is a toxin.

30. A pharmaceutical composition according to claim 26 or claim 27 or claim 28 or claim 29 wherein the ScFv Ab is capable of recognising a 5T4 antigen.

20 31. Use of a ScFv Ab in the preparation of a pharmaceutical composition according to claims 26-30 for the treatment of a condition associated with a DACSM.

25 32. Use of a ScFv Ab capable of recognising a DACSM according to claim 1 in combination with another therapeutically useful agent as defined in claims 27-30 or a nucleotide sequence of interest (NOI) encoding same for the treatment of condition associated with a DACSM.

33. Use of a ScFv Ab according to claims 31 or 32 for *in vivo* imaging and/or for adjuvant treatment of a disease associated with a DACSM.

30

34. Use according to claims 31-33 wherein the disease is cancer.

35. Use of an ScFv Ab to screen for agents that can modulate the DACSM binding specificity of a ScFv Ab wherein the ScFv Ab is an ScFv Ab as defined in claims 1-5 or according to claim 17 or claims 23-24 or is expressed by a nucleotide sequence according to claims 6-12 or a variant, homologue, fragment or derivative thereof.

5

36. A process for diagnosing a disease condition relating to the expression and/or activity of a DACSM in an individual comprising:

10 (i) providing a nucleotide sequence encoding a ScFv Ab as defined in claims 6-12 or an expression product thereof;

(ii) analysing for the binding of the ScFv Ab to a DACSM in a sample derived from the individual;

15 wherein the binding is indicative of the presence of the DACSM in the individual.

37. A method for inducing a therapeutic response in a mammal with a disease condition associated with a DACSM *in vivo* which comprises inoculating the mammal with a ScFv Ab as defined in claims 1-5 or according to claim 17 or claims 23-24 or a vector to direct expression of a nucleotide sequence according to claims 6-12 or a variant, homologue, fragment or derivative thereof in order to induce a therapeutic response to protect said mammal from the disease condition.

20

38. A method according to claim 37 wherein the disease condition is a cancer.

25

39. The use of a ScFv Ab substantially as described herein and with reference to the accompanying Figures.

40. An ScFv for use as a pharmaceutical.

30

ABSTRACT**ANTIBODIES**

- 5 The use of an ScFv Ab (ScFv Ab) capable of recognising a disease associated cell surface molecule (DACSM) in the manufacture of a medicament for the prevention and/or treatment of a disease condition associated with a DACSM is described. The ScFv Ab has therapeutic, diagnostic and prognostic applications.

10

15

20

25

30

SEQUENCE LISTING PART OF THE DESCRIPTION**SEQ ID NO. 1 and SEQ ID No 5**

See Figure 1 (5T4ScFv.1)

5

SEQ ID NO. 3 and SEQ ID No 7

See Figure 2 (B7-1. 5T4ScFv.1)

10 **SEQ ID NO. 4 and SEQ ID No 8**

See Figure 6 (5T4ScFv.1-IgG)

SEQ ID NO. 9 and SEQ ID No 10

15

See Figure 4 (B7-2.5T4.1)

SEQ ID No 11

20 Figure 5 (B7 link ScFv)

SEQ ID NO.12

Figure 7 (ScFv-IgE)

25

SEQ ID NO. 13

Figure 8 (B7-EGF)

30

Figure 1.

1 GAGGTCCAGC TTCAGCAGTC TGGACCTGAC CTGGTGAAGC CTGGGGCTTC
 E V Q L Q Q S G P D L V K P G A S
 51 AGTGAAGATA TCCTGCAAGG CTTCTGGTTA CTCATTCACT GGCTACTACA
 V K I S C K A S G Y S F T G Y Y
 101 TGCACTGGGT GAAGCAGAGC CATGGAAAGA GCCTTGAGTG GATTGGACGT
 M H W V K Q S H G K S L E W I G R
 151 ATTAATCCTA ACAATGGTGT TACTCTCTAC AACCAGAAAT TCAAGGACAA
 I N P N N G V T L Y N Q K F K D K
 201 GGCCATATTA ACTGTAGACA AGTCATCCAC CACAGCCTAC ATGGAGCTCC
 A I L T V D K S S T T A Y M E L
 251 GCAGCCTGAC ATCTGAGGAC TCTGCGGTCT ATTACTGTGC AAGATCTACT
 R S L T S E D S A V Y Y C A R S T
 301 ATGATTACGA ACTATGTTAT GGACTACTGG GGTCAAGTAA CCTCAGTCAC
 M I T N Y V M D Y W G Q V T S V T
 351 CGTCTCCTCA GGTGGTGGTG GGAGCGGTGG TGGCGGCACT GGC GGCGGCG
 V S S G G G G S G G G G T G G G
 401 GATCTAGTAT TGTGATGACC CAGACTCCCA CATTCTGCT TGTTCAGCA
 G S S I V M T Q T P T F L L V S A
 451 GGAGACAGGG TTACCATAAC CTGCAAGGCC AGTCAGAGTG TGAGTAATGA
 G D R V T I T C K A S Q S V S N D
 501 TGTAGDTTGG TACCAACAGA AGCCAGGGCA GTCTCCTACA CTGCTCATAT
 V A W Y Q Q K P G Q S P T L L I
 551 CCTATACATC CAGTCGCTAC GCTGGAGTCC CTGATCGCTT CATTGGCAGT
 S Y T S S R Y A G V P D R F I G S
 601 GGATATGGGA CGGATTTTAC TTTCACCATC AGCACTTTGC AGGCTGAAGA
 G Y G T D F T F T I S T L Q A E D
 651 CCTGGCAGTT TATTTCTGTC AGCAAGATTA TAATTCTCCT CCGACGTTTC
 L A V Y F C Q Q D Y N S P P T F
 701 GTGGAGGCAC CAAGCTGGAA ATCAAACGG
 G G G T K L E I K R

THIS PAGE BLANK (USPTO)

Figure 2

ATGGGCCACA CACGGAGGCA GGGAACATCA CCATCCAAGT GTCCATACCT	50
M G H T R R Q G T S P S K C P Y L	
CAATTTCTTT CAGCTCTTGG TGCTGGCTGG TCTTTCTCAC TTCTGTTTCT	100
N F F Q L L V L A G L S H F C S	
GTGTTATCCA CGTGACCAAG GAAGTGAAAG AAGTGGCAAC GCTGTCCTGT	150
G V I H V T K E V K E V A T L S C	
GGTCACAATG TTTCTGTTGA AGAGCTGGCA CAAACTCGCA TCTACTGGCA	200
G H N V S V E E L A Q T R I Y W Q	
AAAGGAGAAG AAAATGGTGC TGACTATGAT GTCTGGGGAC ATGAATATAT	250
K E K K M V L T M M S G D M N I	
GGCCCGAGTA CAAGAACCGG ACCATCTTTG ATATCACTAA TAACCTCTCC	300
W P E Y K N R T I F D I T N N L S	
ATTGTGATCC TGGCTCTGCG CCCATCTGAC GAGGGCACAT ACGAGTGTGT	350
I V I L A L R P S D E G T Y E C V	
TGTTCTGAAG TATGAAAAAG ACGCTTTCAA GCGGGAACAC CTGGCTGAAG	400
V L K Y E K D A F K R E H L A E	
TGACGTTATC AGTCAAAGCT GACTTCCCTA CACCTAGTAT ATCTGACTTT	450
V T L S V K A D F P T P S I S D F	
GAAATTCCAA CTTCTAATAT TAGAAGGATA ATTTGCTCAA CCTCTGGAGG	500
E I P T S N I R R I I C S T S G G	
TTTTCCAGAG CCTCACCTCT CCTGGTTGGA AAATGGAGAA GAATTAAATG	550
F P E P H L S W L E N G E E L N	
CCATCAACAC AACAGTTTCC CAAGATCCTG AACTGAGCT CTATGCTGTT	600
A I N T T V S Q D P E T E L Y A V	
AGCAGCAAAC TGGATTTCAA TATGACAACC AACCACAGCT TCATGTGTCT	650
S S K L D F N M T T N H S F M C L	
CATCAAGTAT GGACATTTAA GAGTGAATCA GACCTTCAAC TGGAATACAA	700
I K Y G H L R V N Q T F N W N T	

THIS PAGE BLANK (USPTO)

CCAAGCAAGA GCATTTTCCT GATGGAGGCG GGGGATCCGA GGTCCAGCTT	750
T K Q E H F P D G G G G S E V Q L	
CAGCAGTCTG GACCTGACCT GGTGAAGCCT GGGGCTTCAG TGAAGATATC	800
Q Q S G P D L V K P G A S V K I S	
CTGCAAGGCT TCTGGTTACT CATTCACTGG CTACTACATG CACTGGGTGA	850
C K A S G Y S F T G Y Y M H W V	
AGCAGAGCCA TGGAAAGAGC CTTGAGTGGA TTGGACGTAT TAATCCTAAC	900
K Q S H G K S L E W I G R I N P N	
AATGGTGTTA CTCTCTACAA CCAGAAATTC AAGGACAAGG CCATATTAAC	950
N G V T L Y N Q K F K D K A I L T	
TGTAGACAAG TCATCCACCA CAGCCTACAT GGAGCTCCGC AGCCTGACAT	1000
V D K S S T T A Y M E L R S L T	
CTGAGGACTC TGCGGTCTAT TACTGTGCAA GATCTACTAT GATTACGAAC	1050
S E D S A V Y Y C A R S T M I T N	
TATGTTATGG ACTACTGGGG TCAAGTAACC TCAGTCACCG TCTCCTCAGG	1100
Y V M D Y W G Q V T S V T V S S G	
TGGTGGTGGG AGCGGTGGTG GCGGCACTGG CGGCGGCGGA TCTAGTATTG	1150
G G G S G G G G T G G G G S S I	
TGATGACCCA GACTCCCACA TTCCTGCTTG TTTCAGCAGG AGACAGGGTT	1200
V M T Q T P T F L L V S A G D R V	
ACCATAACCT GCAAGGCCAG TCAGAGTGTG AGTAATGATG TAGCTTGGTA	1250
T I T C K A S Q S V S N D V A W Y	
CCAACAGAAG CCAGGGCAGT CTCCTACACT GCTCATATCC TATACATCCA	1300
Q Q K P G Q S P T L L I S Y T S	
GTCGCTACGC TGGAGTCCCT GATCGCTTCA TTGGCAGTGG ATATGGGACG	1350
S R Y A G V P D R F I G S G Y G T	
GATTTCACTT TCACCATCAG CACTTTGCAG GCTGAAGACC TGGCAGTTTA	1400
D F T F T I S T L Q A E D L A V Y	
TTTCTGTCAG CAAGATTATA ATTCTCCTCC GACGTTCCGT GGAGGCACCA	1450
F C Q Q D Y N S P P T F G G G T	

THIS PAGE BLANK (USPTO)

AGCTGGAAAT CAAATAA

K L E I K .

THIS PAGE BLANK (USPTO)

Figure 3a




Figure 3b



THIS PAGE BLANK (USPTO)

Figure 4

1 ATGGGACTGA GTAACATTCT CTTTGTGATG GCCTTCCTGC TCTCTGGTGC
 M G L S N I L F V M A F L L S G A

51 TGCTCCTCTG AAGATTCAAG CTTATTTCAA TGAGACTGCA GACCTGCCAT
 A P L K I Q A Y F N E T A D L P

101 GCCAATTTGC AAACCTCTCAA AACCAAAGCC TGAGTGAGCT AGTAGTATTT
 C Q F A N S Q N Q S L S E L V V F

151 TGGCAGGACC AGGAAAACCTT GGTTCCTGAAT GAGGTATACT TAGGCAAAGA
 W Q D Q E N L V L N E V Y L G K E

201 GAAATTTGAC AGTGTTCATT CCAAGTATAT GGGCCGCACA AGTTTTGATT
 K F D S V H S K Y M G R T S F D

251 CGGACAGTTG GACCCTGAGA CTTCAACAATC TTCAGATCAA GGACAAGGGC
 S D S W T L R L H N L Q I K D K G

301 TTGTATCAAT GTATCATCCA TCACAAAAAG CCCACAGGAA TGATTCGCAT
 L Y Q C I I H H K K P T G M I R I

351 CCACCAGATG AATTCTGAAC TGTCAGTGCT TGCTAACTTC AGTCAACCTG
 H Q M N S E L S V L A N F S Q P

401 AAATAGTACC AATTTCTAAT ATAACAGAAA ATGTGTACAT AAATTTGACC
 E I V P I S N I T E N V Y I N L T

451 TGCTCATCTA TACACGGTTA CCCAGAACCT AAGAAGATGA GTGTTTTGCT
 C S S I H G Y P E P K K M S V L L

501 AAGAACCAAG AATTCAACTA TCGAGTATGA TGGTATTATG CAGAAATCTC
 R T K N S T I E Y D G I M Q K S

551 AAGATAATGT CACAGAACTG TACGACGTTT CCATCAGCTT GTCTGTTTCA
 Q D N V T E L Y D V S I S L S V S

601 TTCCCTGATG TTACGAGCAA TATGACCATC TTCTGTATTC TGGAAACTGA
 F P D V T S N M T I F C I L E T D

651 CAAGACGCGG CTTTTATCTT CACCTTTCTC TATAGAGCTT GAGGACCCTC
 K T R L L S S P F S I E L E D P

THIS PAGE BLANK (USPTO)

AGCCTCCCCC AGACCACATT CCTGGAGGCG GGGGATCC
Q P P P D H I P G G G G S

THIS PAGE BLANK (USPTO)

Figure 5

atggccttgca	attgtcagtt	gatgcaggat	acacccactcc	tcaagtttcc	atgtccaagg	60
ctcattcttc	tctttgtgct	gctgattcgt	ctttcacaag	tgtcttcaga	tgttgatgaa	120
caactgtcca	agtcagtgaa	agataaggta	ttgctgcctt	gccgttacaa	ctctccgcat	180
gaagatgagt	ctgaagaccg	aatctactgg	caaaaacatg	acaaagtgg	gctgtctgtc	240
attgctggga	aactaaaagt	gtggcccag	tataagaacc	ggactttata	tgacaacact	300
acctactctc	ttatcatcct	gggcctgggc	ctttcagacc	ggggcacata	cagctgtgtc	360
gttcaaaaga	aggaaagagg	aacgtatgaa	gttaaact	tggctttagt	aaagttgtcc	420
atcaaagctg	acttctctac	ccccaacata	actgagtctg	gaaacccatc	tgacagacact	480
aaaaggatta	cctgctttgc	ttccgggggt	ttcccaaagc	ctcgcttctc	ttggttgga	540
aatggaagag	aattacctgg	catcaatacg	acaatttccc	aggatcctga	atctgaattg	600
tacaccatta	gtagccaact	agatttcaat	acgactcgca	accacacccat	taagtgtctc	660
attaaatatg	gagatgctca	cgtgtcagag	gacttcacct	gggaaaaacc	cccagaagac	720
cctcctgata	gcaagcccgg	gggtgggtgg	agcgggtggg	gcggcagtg	cggcggcgga	780
actagtgagg	tccagcttca	gcagtctgga	cctgacctgg	tgaagcctgg	ggcttcagtg	840
aagatatcct	gcaaggcttc	tggttactca	ttcactggct	actacatgca	ctgggtgaag	900
cagagccatg	gaaagagcct	tgagtggatt	ggacgtatta	atcctaacaa	tggtgttact	960
ctctacaacc	agaaattcaa	ggacaaggcc	atattaactg	tagacaagtc	atccaccaca	1020
gcctacatgg	agctccgcag	cctgacatct	gaggactctg	cggtctatta	ctgtgcaaga	1080
tctactatga	ttacgaacta	tggttatggac	tactgggggc	aagtaacttc	agtcaccgtc	1140
tcttcagggtg	gtgggtgggag	cggtgggtggc	ggcactggcg	gcggcggatc	tagtattgtg	1200
atgaccacaga	ctcccacatt	cctgcttggt	tcagcaggag	acagggttac	cataacctgc	1260
aaggccagtc	agagtgtgag	taatgatgta	gcttgggtacc	aacagaagcc	agggcagttc	1320
cctacactgc	tcataatccta	tacatccagt	cgctacgctg	gagtccctga	tcgcttcatt	1380
ggcagtggtat	atgggacgga	tttcactttc	accatcagca	ctttgcaggc	tgaagacctg	1440
gcagttttatt	tctgtcagca	agattataat	tctcctccga	cgttcggtgg	aggcaccaag	1500
ctggaaatca	aacggtaa					1518

THIS PAGE BLANK (USPTO)

Figure 6

Leader / 5T4 scFv / HlgG DNA and deduced protein sequence

CTCGAGCCACCATGGGATGGAGCTGTATCATCCTCTTCTTGGTAGCAACAGCTACAGGTGTCCACTCCGAGGTCCAGCTG
 M G W S C I I L F L V A T A T G V H S E V Q L
 CAGCAGTCTGGACCTGACCTGGTGAAGCCTGGGGCTTCAGTGAAGATATCCTGCAAGGCTTCTGGTTACTCATTCACTGG
 Q Q S G P D L V K P G A S V K I S C K A S G Y S F T
 CTACTACATGCACTGGGTGAAGCAGAGCCATGGAAAGAGCCTTGAGTGGATTGGACGTATTAATCCTAACAATGGTGTTA
 G Y Y M H W V K Q S H G K S L E W I G R I N P N N G V
 CTCTCTACAACCAGAAATTCAAGGACAAGGCCATATTAAGTGTAGACAAGTCATCCACCACAGCCTACATGGAGCTCCGC
 T L Y N Q K F K D K A I L T V D K S S T T A Y M E L R
 AGCCTGACATCTGAGGACTCTGCGGTCTATTACTGTGCAAGATCTACTATGATTACGAATATGTTATGGACTACTGGGG
 S L T S E D S A V Y Y C A R S T M I T N Y V M D Y W
 TCAAGTAATTCAGTCACCGTCTCTTCAGGTGGTGGTGGGAGCGGTGGTGGCGGCACTGGCGGCGGCGGATCTAGTATTG
 G Q V T S V T V S S G G G G S G G G G T G G G G S S I
 TGATGACCCAGACTCCACATTCTGCTTGTTCAGCAGGAGACAGGGTTACCATAACCTGCAAGGCCAGTCAGAGTGTG
 V M T Q T P T F L L V S A G D R V T I T C K A S Q S V
 AGTAATGATGTAGCTTGGTACCAACAGAAGCCAGGCGAGTCTCTACACTGCTCATATCCTATACATCCAGTCGCTACGC
 S N D V A W Y Q Q K P G Q S P T L L I S Y T S S R Y
 TGGAGTCCCTGATCGCTTCATTGGCAGTGGATATGGGACGGATTTCACTTTCACCATCAGCACTTTGCAGGCTGAAGACC
 A G V P D R F I G S G Y G T D F T F T I S T L Q A E D
 TGGCAGTTTATTTCTGTGACGAAGATTATAATTCTCTCCGACGTTCCGGTGGAGGCACCAAGCTTGAAATCAAACGGGCC
 L A V Y F C Q Q D Y N S P P T F G G G T K L E I K R A
 TCCACCAAGGGCCCATCGGTCTTCCCCCTGGCACCCCTCCTCCAAGAGCACCTCTGGGGGCACAGCGGCCCTGGGCTGCCT
 S T K G P S V F P L A P S S K S T S G G T A A L G C
 GGTCAGGACTACTTCCCCGAACCGGTGACGGTGTCTGGAAGTCAAGCGCCCTGACCAGCGCGTGCACACCTTCCCGG
 L V K D Y F P E P V T V S W N S G A L T S G V H T F P
 CTGTCCTACAGTCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCAGCTTGGGCACCCAGACCTAC
 A V L Q S S G L Y S L S S V V T V P S S S L G T Q T Y
 ATCTGCAACGTGAATCACAAGCCAGCAACACCAAGGTGGACAAGAAAGTTGAGCCCAAATCTTGTGACAAAACCTCACAC
 I C N V N H K P S N T K V D K K V E P K S C D K T H
 ATGCCCACCGTGCCAGCACCTGAACTCCTGGGGGGACCGTCAGTCTTCTCTTCCCCCAAACCAAGGACACCCTCA
 T C P P C P A P E L L G G P S V F L F P P K P K D T L
 TGATCTCCCGGACCCCTGAGGTACATGCGTGGTGGTGGACGTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTAC
 M I S R T P E V T C V V V D V S H E D P E V K F N W Y
 GTGGACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGAGGAGCAGTACAACAGCACGTACCGTGTGGTCAAGCT
 V D G V E V H N A K T K P R E E Q Y N S T Y R V V S
 CCTCACCGTCTGCACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGAAGGTCTCCAACAAAGCCCTCCCAGCCCCCA
 V L T V L H Q D W L N G K E Y K C K V S N K A L P A P
 TCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCATCCCGGGATGAGCTG
 I E K T I S K A K G Q P R E P Q V Y T L P P S R D E M
 ACCAAGAACCAGGTGAGCTGACCTGCCTGGTCAAAGGCTTCTATCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGG
 T K N Q V S L T C L V K G F Y P S D I A V E W E S N
 GCAGCCGGAGAACAACATAAGACCACGCTCCCGTGGTGGACTCCGACGGCTCCTTCTCTCTATAGCAAGCTCACCG
 G Q P E N N Y K T T P P V L D S D G S F F L Y S K L T
 TGGACAAGAGCAGGTGGCAGCAGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACGCAG
 V D K S R W Q Q G N V F S C S V M H E A L H N H Y T Q
 AAGAGCCTCTCCCTGTCCCCGGGTAAATGACTCGAG
 K S L S L S P G K .

THIS PAGE BLANK (USPTO)

Figure 7

ctcgagccac	catgggatgg	agctgtatca	tcctcttctt	ggtagcaaca	gctacagggtg	60
tccactccga	gggccagctg	cagcagctctg	gacctgacct	ggggaagcct	ggggcttcag	120
tgaagatatc	ctgcaaggct	tctgggttact	cattcactgg	ctactacatg	cactgggtga	180
agcagagcca	tggaaagagc	cttgagtggg	ttggacgtat	taatcctaac	aatgggtgta	240
ctctctacaa	ccagaaattc	aaggacaagg	ccatattaac	tgtagacaag	tcatccacca	300
cagcctacat	ggagctccgc	agcctgacat	ctgaggactc	tgcgggtctat	tactgtgcaa	360
gatctactat	gattacgaac	tatgttatgg	actactgggg	tcaagtaact	tcagtcaccg	420
tctcttcagg	tgggtgggtgg	agcgggtggg	gcggcactgg	cggcgggcga	tctagtattg	480
tgatgaccca	gactcccaca	ttcctgcttg	tttcagcagg	agacagggtt	accataacct	540
gcaaggccag	tcagagtgtg	agtaatgatg	tagcttggtg	ccaacagaag	ccaggggcagt	600
ctcctacact	gctcatatcc	tatacatcca	gtcgctacgc	tggagtccct	gactcgcttca	660
ttggcagtg	atatgggacg	gatttcactt	tcaccatcag	cactttgcag	gctgaagacc	720
tggcagttta	tttctgtcag	caagattata	attctcctcc	gacgttcggt	ggaggcacca	780
agcttgaaat	caaacggggc	tccacacaga	gccatccgt	cttccccctg	acccgctgct	840
gcaaaaacat	tccttccaat	gccacctccg	tgactctggg	ctgcctggcc	acgggctact	900
tcccggagcc	ggtgatgggtg	acctgggaca	caggctccct	caacgggaca	actatgacct	960
taccagccac	cacctcacg	ctctctgggtc	actatgccac	catcagcttg	ctgaccgtct	1020
cgggtgcgtg	ggccaagcag	atgttcacct	gccgtgtggc	acacactcca	tcgtccacag	1080
actgggtcga	caacaaaacc	ttcagcgtct	gctccaggga	cttcacccccg	cccaccgtga	1140
agatcttaca	gtcgtcctgc	gacggcggcg	ggcacttccc	cccgaccatc	cagctcctgt	1200
gcctcgtctc	tgggtacacc	ccagggaacta	tcaacatcac	ctggctggag	gacgggcagg	1260
tcattggacgt	ggacttgctc	accgcctcta	ccacgcagga	gggtgagctg	gcctccacac	1320
aaagcgagct	cacctcagc	cagaagcact	ggctgtcaga	ccgcacctac	acctgccagg	1380
tcacctatca	aggtcacacc	tttgaggaca	gcaccaagaa	gtgtgcagat	tccaaccgga	1440
gaggggtgag	cgctaccta	agccggccca	gcccgttcga	cctgttcac	cgcaagtcgc	1500
ccacgatcac	ctgtctgggtg	gtggacctgg	cacccagcaa	ggggaccgtg	aacctgacct	1560
ggtcccgggc	cagtgggaag	cctgtgaacc	actccaccag	aaaggaggag	aagcagcgca	1620
atggcacgtt	aaccgtcacg	tccaccctgc	cgggtgggcac	ccgagactgg	atcgaggggg	1680
agacctacca	gtgcagggtg	acccaccccc	acctgcccag	ggccctcatg	cgggtccacga	1740
ccaagaccag	cgccccgcgt	gctgccccgg	aagtctatgc	gtttgcgacg	ccggagtggc	1800
cggggagccg	ggacaagcgc	accctcgctc	gcctgatcca	gaacttcacg	cctgaggaca	1860
tctcggtgca	gtggctgcac	aacgaggtgc	agctcccggg	cgcccggcac	agcacgacgc	1920
agccccgcaa	gaccaagggc	tccggcttct	tcgtcttcag	ccgcctggag	gtgaccaggg	1980
ccgaatggga	gcagaaagat	gagttcatct	gccgtgcagt	ccatgaggca	gcgagcccct	2040
cacagaccgt	ccagcgagcg	gtgtctgtaa	atcccggtaa	atgagagctc		2090

THIS PAGE BLANK (USPTO)

Figure 8

```


atggcttgca attgtcagtt gatgcaggat acaccactcc tcaagtttcc atgtccaagg 60
ctcattcttc tctttgtgct gctgattcgt ctttcacaag tgtcttcaga tgttgatgaa 120
caactgtcca agtcagtgaag agataaggta ttgctgcctt gccgttacaa ctctccgcat 180
gaagatgagt ctgaagaccg aatctactgg caaaaacatg acaaagtggg gctgtctgtc 240
attgctggga aactaaaagt gtggcccagag tataagaacc ggactttata tgacaacact 300
acctactctc ttatcatcct gggcctgggc ctttcagacc ggggcacata cagctgtgtc 360
gttcaaaaaga aggaaagagg aacgtatgaa gttaaactact tggctttagt aaagtgtgcc 420
atcaaagctg acttctctac cccaacata actgagtctg gaaacccatc tgcagacact 480
aaaaggatta cctgctttgc ttccgggggt ttcccaaagc ctgcttctc ttggttgaa 540
aatggaagag aattacctgg catcaatacg acaatttccc aggatcctga atctgaattg 600
tacaccatta gtagccaact agatttcaat acgactcgca accacaccat taagtgtctc 660
attaaatatg gagatgctca cgtgtcagag gacttcacct gggaaaaacc cccagaagac 720
cctcctgata gcaagcccgg ggtggtggg agcgggtggtg gcggcagtgg cggcggcgga 780
actagtaata gtgactctga atgtcccctg tcccacgatg ggtactgcct ccatgatggg 840
gtgtgcatgt atattgaagc attggacaag tatgcatgca actgtgttgt tggctacatc 900
ggggagcgat gtcagtaccg agacctgaag tgggtgggaac tgcgc 945

```

THIS PAGE BLANK (USPTO)

Figure 9


CT26-neo Transfectants



THIS PAGE BLANK (USPTO)

Figure 10


CT26-h5T4 Transfectants



THIS PAGE BLANK (USPTO)

Figure 11


B16-neoTumour Growth



THIS PAGE BLANK (USPTO)


Figure 12

B16-h5T4 Tumour Growth




THIS PAGE BLANK (USPTO)

Figure 13




THIS PAGE BLANK (USPTO)

Figure 14



THIS PAGE BLANK (USPTO)


Figure 15




THIS PAGE BLANK (USPTO)

Figure 16

A: A9-5T4



B: A9-neo



THIS PAGE BLANK (USPTO)

Figure 17



Cell Lines & E:T Ratio

Fusion Protein

THIS PAGE BLANK (USPTO)